


# CLINICAL EEG

ELECTROENCEPHALOGRAPHY



## Author Index

Vol. 18, 1987

## Subject Index

Vol. 14, 1983

Vol. 15, 1984

Vol. 16, 1985

Vol. 17, 1986

Vol. 18, 1987

# Index of Authors

---

- Aguglia, Umberto, 12, 147  
 Amit, R., 169  
 Ammoumi, A.A., 233  
 Asokan, G., 201  
 Awerbuch, Gavin I., 38  
 Balachandran, Suppiah, 20  
 Baralt, Tulio, 187  
 Beadle, Beverly, 74  
 Becker, Jacqueline, 93  
 Bradshaw, C. Marshall, 164  
 Buchsbaum, Monte S., 61  
 Calderon, Froylan E., 187  
 Caldwell, Diane L., 26  
 Carney, Andrew L., No. 2, IX  
 Carra, Joseph P., 15  
 Chesser, Michael Z., 227  
 Chu, Nai-Shin, 159  
 Clarke, M., 68  
 Congbalay, Dean R., 81, 136  
 Corsaro, Carole J., 124  
 Dadmehr, Nahid, 34, 81, 136  
 Dang, Minh-Tam, 20  
 Drake, Jr., Miles E., 34, 81, 136  
 Eden, O.B., 68  
 El-Mallakh, Rif S., 2  
 Erkulwater, Samard, 211  
 Espinosa Tanguna, Roberto, 187  
 Farnarier, G., 147  
 Fiorelli, Sergio, 85  
 Fisher, R.S., 191  
 Franco Delgadillo, Juan, 89  
 Frazier, E., 103  
 Fukushima, Yutaka, 114  
 Furune, S., 120  
 Gamboa Marrufo,  
     Jose Domingo, 187  
 Ganji, S., 103, 180  
 Gastaut, Henri, 12  
 Ghaly, Ramsis F., 218  
 Gibbs, Erna L., 10  
 Gibbs, Frederic A., 10  
 Gomez, M., 147  
 Greiffenstein, Manfred F., 26  
 Halabe, Jose, 85  
 Happel, Lee, 180  
 Harmony, Thalia, 93  
 Harrington, Donna, 20  
 Hays, Amy L., 15  
 Hughes, John R., 163, 218  
 Jacome, Daniel E., 41  
 Janati, Abdorasool, 20, 74, 227  
 Jeub, Robert P., 164  
 Kellams, Jeffrey J., 124  
 Kidwai, Shahab, 20  
 King, Suzanne D., 26  
 Kubota, Shuji, 114  
 Landazuri, Ana Maritza, 93  
 Laudadio, Silvia, 12  
 Lee, Sunghoon, 61  
 Leonberg, Jr., Stanley C.,  
     No. 2, IX  
 Lifshitz, Alberto, 85  
 Lopez, Moises, 85  
 Maehara, M., 120  
 Malagon Valdez, Jorge, 89  
 Marino, Steven, 180  
 Marosi, Erzsebet, 93  
 McInnes, A., 68  
 Medina, Eloy, 85  
 Miller, Marvin J., 124  
 Milstein, Victor, 124  
 Nazarian, Sarkis M., 227  
 Negoro, T., 120  
 Niedermeyer, E., 191, 201  
 Nigro, M., 142  
 Nomura, K., 120  
 Nowack, William J., 74  
 O'Hare, A.E., 68  
 Olmos Garcia de Alba,  
     Graciela, 89, 187  
 Osuna, Guadalupe, 85  
 Pakalnis, Ann, 34, 136  
 Pareja, J., 201  
 Pena, Yolanda M., 2  
 Peters, G., 103  
 Podurgiel, Bernard J., 2  
 Potts, Robert E., 227  
 Quattrone, A., 147  
 Ramos Peek, Jaime, 89  
 Reilly, Edward L., 167  
 Rengifo Ramos, Oscar, 187  
 Rey, M., 147  
 Rivera Olmos, Victor M., 187  
 Roccaforte, Peter, 218  
 Rodin, E., 142  
 Saito, Fumio, 114  
 Sandford, Sandra L., 15  
 Sciabassi, Robert J., 15  
 Shanley, John D., 2  
 Shulka, D., 233  
 Small, Iver F., 124  
 Small, Joyce G., 124  
 Soll, Robert W., 163  
 Stone, James L., 218  
 Struve, Frederick A., 46, 173  
 Subramanian,  
     Kodanallur S., 218  
 Synek, B.J.L., 54  
 Synek, V.M., 54  
 Takahashi, I., 120  
 Tarter, Ralph E., 15  
 Tilton, Ann C., 180  
 Tinuper, Paolo, 12, 147  
 Toggia, Joseph U., 165  
 Van Thiel, David H., 15  
 Velasco, Marcos, 93  
 Verma, Narayan P., 26, 38  
 Verma, Neelam, 26  
 Watanabe, K., 120  
 Yamamoto, N., 120

# Index of Subjects

- Abdominal epilepsy** and, valproic acid treatment in children, 17:14-20
- Abscess, brain**, and, coma in children, 14:207-213
- Absence seizures** and, familial photosensitive epilepsy, 15:47-52
- Lennox-Gastaut syndrome, 17:117-126
  - mentally retarded, seizure management, 14:164-172
  - mesothalamic reticular stimulation, 17:52-60
  - natural history of hypsarrhythmia 16:128-130
  - status appearing on eye-closure, 16:111-118
  - status, termination and suppression of interictal bursts with phenytoin, case report, 17:173-175
  - treatment, epileptic syndromes in childhood, book review, 17:No. 3, VII
  - valproic acid treatment in children, 17:14-20
- Academic achievement**, and, children with learning disabilities, 18:93-101
- Acronyms**, and use in EEG, 15:32-39
- ACTH** and, hypsarrhythmia, diminishing incidence, 16:178-182
- infantile spasms and hypsarrhythmia, early success and ongoing benefits of Sorel's treatment, 16:120-121
  - Lennox-Gastaut syndrome, 17:117-126
  - West's syndrome, treatment, 15:140-144; 16:120-121, 122-127
- Acupuncture**, and effect on somatosensory evoked potentials, 14:53-56
- Adult EEG** and, centenarians, 17:146-151
- episodic anterior drowsy theta, 17:135-138
  - herpes simplex virus encephalitis, survival in an elderly man, with periodic sharp complexes, 18:2-9
  - mid-line epileptiform activity, 16:54-59
  - paroxysmal hyperventilation responses, 17:61-65
  - patients with fever, 18:85-88
  - POSTS, age-dependent incidence, 16:11-15
  - psychogeriatric diagnosis, photic driving, 14:78-81
  - stuttering and seizures, 17:142-145
  - theta pattern coma in younger adults, 18:54-60
- Affective disorders** and, pattern reversal evoked potentials, 15:178-183
- tardive dyskinesia, 15:222-225
- Age factor** and, alcohol abuse with pattern reversal VEP, 15:102-109
- centenarians, observations of the EEG in 7 cases, 17:146-151
  - episodic anterior drowsy theta, 17:135-138
  - fever, EEG in adult patients, 18:85-88
  - geriatric disease with photic driving, 14:78-81
  - hyperventilation responses, 17:61-65
  - hypsarrhythmia, infection as a possible etiology, 17:78-81
  - hypsarrhythmia, natural history, 16:128-130
  - Lennox-Gastaut syndrome, 17:117-126
  - mu rhythm, 17:44-50
  - pattern reversal VEP, latency changes, 18:159-162
  - POSTS in geriatric EEG, 16:11-15
  - rhythmical temporal theta, 15: 145-150
  - small sharp spikes, 15:208-213
  - theta pattern coma occurring in younger adults, 18:54-60
  - (See also Adult EEG, Children and Neonates)
- Akinetic seizures** and, Lennox-Gastaut syndrome, 17:117-126
- Alcohol** and, toxicity in pediatric cases of coma, 14:207-213
- Alcoholism** and, EEG pattern in delirium tremens, 16:30-32
- pattern reversal visual evoked potentials, effect of age, 15:102-109
  - short latency brainstem and somatosensory evoked potentials, 14:8-16
  - spindle coma in benzodiazepine toxicity, case report, 16:213-218
  - visual evoked potentials, 14:17-19
- Alpha activity** and, bipolar vs. monopolar recording, 15:1-21, 71-82
- centenarians, 17:146-151
  - cerebral infarction, 18:20-25
  - children with end-stage renal disease before and after haemodialysis, 17:195-202
  - coma, 15:32-39, 116-121, 167-172; 17:82-88, 187-194; 18:54-60, 74-80, 103-113
  - delirium tremens, 16:30-32
  - EEG topography in porencephaly and arachnoid cyst, 16:21-29
  - ECT, analysis of spectral energy, 17:66-77
  - ECT with EEG spectral energy distributions and methohexital, Innovar and ketamine anesthetics, 17:203-215
  - evoked potential and clinical studies in alpha coma, 18:103-113
  - neonatal meningitis, with theta, 14:67-77
  - neonatal status epilepticus, 15:125-131
  - normal subjects with lorazepam and methylphenidate, effect on event-related potentials, 17:176-180
  - parkinsonian patients on levodopa, 14:82-85
  - periodic activity, 15:32-39
  - post-traumatic syndrome, 15:214-221
  - psychiatric inpatients with antipsychotic-induced tardive dyskinesia, 17:30-35

- schizophrenia, brain surface activity mapping, 18:124-135
- spectral analysis of interictal EEG in epilepsy, 16:94-97
- topographic mapping, artifacts, 18:61-67
- vascular brain stem lesions, 14:35-46
- Alpha coma** and, alpha rhythm and alpha-like coma, 15:167-172
  - anoxic encephalopathy, coexistence with periodic EEG pattern, 17:187-194
  - coexisting with theta pattern coma and spindle coma, 18:74-80
  - evoked potential and clinical studies, 18:103-113
  - periodic EEG activity, 15:32-39
  - suppression-burst pattern with generalized epileptiform discharges, 17:82-88
  - theta pattern coma, a variant of alpha pattern coma, 15:116-121
  - theta pattern coma prognostic significance, 18:54-60
- Alzheimer disease** and, electrophysiologic validation of dementias, cortical and sub-cortical, 18:26-33
  - event related evoked potentials, 16:48-53
  - periodic EEG activity, 15:32-39
  - pre-senile atrophy, with EEG changes, 18:164
- Ambulatory EEG** and, community hospital prospective study with analysis of patient outcome, 16:161-166
  - monitoring of all night sleep, 16:16-20
  - pediatrics, 24 hour monitoring, 15:164-166
  - post-traumatic syndrome, 15:214-221
  - whiplash injury, 18:41-45
- American Medical EEG Association (AMEEGA)** and, EEG instrument standards, 15:No. 3, VI
- Anatomical placement of EEG electrodes**, and universal APEEGE system, 15:1-21
- Anectine**, and attenuation of EMG with electro-cerebral inactivity, 16:72-76
- Anesthesia** and, ECT with EEG spectral energy distributions with methohexital, Innovar and ketamine, 17:203-215
  - electrocorticograms, 15:83-91
  - spinal cord lesions, effect on SEP, 14:142-151, 173-185
- Animal model** and, spinal cord lesions, effect on SEP, 14:142-151, 173-185
- Anoxia** and, coexistence of ectopic rhythms and periodic EEG pattern, 17:187-194
  - drowning survival in children with EP as predictors of outcome, 16:183-191
  - suppression-burst pattern, epileptiform discharges and alpha-theta pattern coma, 17:82-88
  - theta pattern coma occurring in younger adults, 18:54-60
  - (See also Apnea and Hypoxia)
- Anterior bradyrhythmia**, and frontal inter-mittent rhythmical delta activity, significance of two patterns, 16:1-10
- Anterior temporal localization**, and, electrode placement systems 15:1-21
  - small sharp spikes, possible relationship to epilepsy, 18:114-119
  - (See also Temporal)
- Antiepileptic medication** and absence status, 16:111-118
  - ambulatory EEG study, 16:161-166
  - computerized spectral analysis of interictal EEG in epilepsy, 16:94-97
  - ECT, 17:203-215
  - familial photosensitive epilepsy, 15:47-52
  - interictal EEG prediction of response to monotherapy, 16:131-135
  - Lennox-Gastaut syndrome, 17:117-126
  - management in institutionalized mentally retarded epileptics, 14:164-172
  - neonatal seizures, 15:133-139, 197-201
  - phenytoin infusion in seizure patients, 15:173-177
  - phenytoin termination of absence status and suppression of interictal bursts, 17:173-175
  - serum levels, effect of valproic acid on diphenylhydantoin, 16:104-110
  - sleep and waking disturbances in epilepsy, 18:136-141
  - status epilepticus, neonatal, 15:197-201
  - stuttering and seizures in adults, 17:142-145
  - valproic acid in children, 17:14-20
- Antisocial personality**, and brainstem auditory evoked potential, 16:91-93
  - (See also Behavior and Psychiatric disorders)
- APEEGE**, anatomical placement of EEG electrodes system, 15:1-21
- Apnea** and, muscle activity in the legs in narcolepsy and obstructive sleep apnea, 17:181-186
  - neonatal seizures, 14:67-77; 15:133-139, 193-196
  - reactive burst suppression in infants, 14:138-141
  - sleep disturbances in mentally retarded infants, 14:90-95
  - sleep disturbances in narcolepsy, 14:130-134; 17:181-186
- Arachnoid cyst**, and EEG topography, 16:21-29
- Arousal** and, anterior bradyrhythmia, compared to FIRDA, 16:1-10
  - BAEP in antisocial personality, 16:91-93
  - muscle activity in the legs in narcoleptics and obstructive sleep apnea patients, 17:181-186
  - seizures with pain, role of reticular activating system, 16:195-201
  - unusual patterns in children, 15:92-96
- Arteriovenous malformation** and, FIRDA in

- pial-dural arteriovenous malformation, 18:227-232
- (See also Vascular disease)
- Artifact** and, alcoholics with epilepsy, 14:8-16
- burst suppression pattern, 17:162-168
- delirium tremens, 16:30-32
- EMG, failure of Pavulon to provide attenuation in electrocerebral inactivity, 16:72-76
- lightning, 17:105-109
- management of infants and children for EEG, 15:202-207
- tongue-movement, 14:57-59
- topographic mapping of EEG, 18:61-67
- Asphyxia**, and reactive burst suppression in infants, 14:138-141
- (See also Hypoxia)
- Astatic seizures** and, Lennox-Gastaut syndrome, 17:117-126
- valproic acid treatment in children, 17:14-20
- Asymmetry, EEG**, and multiple sclerosis, 18:38-40
- neurological disorders, diagnosis, 14:116-129
- normal and abnormal crossed asymmetry in monocular visual evoked potentials, 16:77-82
- spindles awake with glioblastoma, 17:1-5
- tuberous sclerosis, 16:219-224
- Asynchrony**, and congenital hydrocephalus, 15:22-31
- Atonic drop attacks**, and Lennox-Gastaut syndrome, 17:117-126; 18:191-200
- Atrophy, cerebral**, and Alzheimer's disease, pre-senile, with EEG changes, 18:164
- epilepsia partialis continua, 16:45-47
- mid-line epileptiform activity in adult EEGs, 16:54-59
- (See also Alzheimer disease)
- Auditory evoked potentials** and, alcoholics, short latency BAER, 14:8-19
- alpha coma, clinical and EP studies, 18:103-113
- Alzheimer disease, 16:48-53
- antisocial personality, 16:91-93
- brain-injured patients with rage attacks and self-injurious behavior, 17:169-172
- brainstem death, 16:63-71
- dementias, cortical and subcortical, 18:26-33
- dizziness, 15:110-115
- drowning outcome, predictors of survival in young children, 16:183-191
- epilepsy; long-latency auditory event-related potentials, 17:10-13
- FIRDA, 14:135-137
- hearing loss, 15:159-163
- hepatic injury status, 18:15-19
- hydrocephalus, 15:22-31
- hypsarhythmia-infantile spasms in near-drowning, 18:180-186
- Lafora disease, 16:202-207
- liver transplantation, 16:192-194
- modified auditory brainstem responses (MABR), Part I - rationale and normative study, 18:218-226
- neonatal meningitis, 14:67-77
- non-convulsive status epilepticus, 16:33-38
- normal subjects, 17:176-180
- sensori-neural hearing loss, 15:159-163
- surgical monitoring, pathological significance of reduced ABR P1-P5 interwave interval, 16:83-87
- Automatisms**, and benign infantile epilepsies, 16:39-44
- complex partial status epilepticus, 14:225-232
- occipital evoked spike epilepsy in childhood, 14:221-224
- Bascale pattern**, and neonatal status epilepticus, 15:197-201
- Behavior disorder** and, BAEP in antisocial personality, 16:91-93
- BAEP in brain-injured patients with rage attacks and self-injurious behavior, 17:169-172
- children with end stage renal disease before and after haemodialysis, 17:195-202
- computerized spectral analysis of interictal EEG in epilepsy, 16:94-97
- electrophysiological tests and post-concussion syndromes, 18:165-167
- epilepsy, success of EEG in confirmation, 16:98-103
- 14 and 6 positive spikes, 15:92-96
- (See also Psychiatric disorder)
- Benign focal epilepsy of childhood**, and individual and intrafamilial multifocality of spikes, 18:169-172
- infantile epilepsies with evoked spikes, 16:39-44
- parietal epilepsy with evoked and spontaneous spikes in siblings, 17:159-161
- Benzodiazepines** and, toxicity with spindle coma, case report, 16:213-218
- Benzotropine** and, computerized EEG profiles in treatment resistant schizophrenia, 18:124-135
- Beta activity** and, Alzheimer's disease, 18:164
- centenarians, 17:146-151
- cerebellar syndrome, cerebral electrical field mapping, 18:142-146
- children with end stage renal disease before and after haemodialysis, 17:195-202
- clinical correlates of very fast activity, 15:237-241
- coma, 14:207-213; 15:32-39, 167-172
- delirium tremens, 16:30-32
- development in children, EEG-topographical analysis, 18:120-123
- electroconvulsive therapy, 17:66-77, 203-215

- grand mal discharges, 15:155-158
- normal patterns in children, 15:92-96
- normal subjects, effect of lorazepam and methylphenidate on event-related potentials and power spectral analysis, 17:176-180
- post-traumatic syndrome, 15:214-221
- schizophrenia, brain surface activity mapping, 18:124-135
- tardive dyskinesia, 17:30-35
- topographic mapping, artifacts, 18:61-67
- whiplash injury, 18:41-45
- BETS**, benign epileptiform transients of sleep further data on a controversial pattern, 15:208-213
  - (See also Small sharp spikes)
- BIPLEDs**, and cerebral fat embolism, 14:27-34
  - (See also PLEDs)
- Bipolar recording** and, APEEGE monopolar system, 15:1-21
  - historical notes, 14: No. 2, V, No. 3, VII, VIII
  - Neo-EEG monopolar system, 15:71-82
- Birth Injury**, (See Perinatal injury)
- Blindness**, (See Visual disorder)
- Blood levels** and barbiturates in children with coma after drowning, 16:183-191
  - valproic acid effect on serum levels of diphenylhydantoin, 16:104-110
- Book Review**, Epilepsy, a Medical Handbook for Physicians, Nurses, Teachers, Parents, 16:No.4, VII
  - Epileptic Syndromes in Infancy, Childhood and Adolescence, 17: No. 3, VII
  - Evoked Potential Monitoring in the Operating Room, 18:No. 2, XII
  - Topographic Mapping of Brain Electrical Activity, 18:No. 1, VIII
- BORTT** (bursts of rhythmical temporal theta), a distinctive EEG entity, 15:145-150
- Brain death**, (See Death)
- Brain electrical activity mapping (BEAM)** and, artifacts, 18:61-67
  - cerebellar syndrome, 18:142-146
  - development of beta activities in children, 18:120-123
  - editorial, advanced methods for EEG recording and analysis, 18:No. 2, IX
  - haloperidol, chlorpromazine, clozapine and placebo in treatment resistant schizophrenia, 18:124-135
  - porencephaly and arachnoid cyst, 16:21-29
- Brainstem auditory evoked responses (BAER)** (see Auditory evoked potentials)
- Brainstem disorder** and, death, 15:No. 1,V,VI
  - dizziness with EP, 15:110-115
  - rage attacks in brain-injured patients, 17:169-172
  - sensori-neural hearing loss, with auditory nerve BER, 15:159-163
  - theta coma, 15:116-121
  - vascular lesions, disturbances of sleep-waking cycle, 14:35-46
- (See also Auditory evoked potentials)
- Bromocriptin**, and therapy effect on the EEG in chronic hepatic encephalopathy, 14:86-89
- Burst suppression** (see Suppression-burst pattern)
- Bursts of rhythmical temporal theta (BORTT)**, a distinctive EEG entity, 15:145-150
- Calcification** and, tuberous sclerosis long term follow-up, 16:219-224
- Carbamazepine** and, hypnogenic paroxysmal dystonia, 16:136-142
  - interictal EEG prediction of response to anti-epileptic drug monotherapy, 16:131-135
  - seizure management in mentally retarded, 14:164-172
  - sleep and waking disturbances in epilepsy, 18:136-141
- Cardiac disorder** and, alpha coma, 15:167-172
  - cardiorespiratory arrest and theta pattern coma occurring in younger adults, 18:54-60
  - dizziness, 15:110-115
  - (See also Vascular disease)
- Carotid endarterectomy**, and assessment of variability of early scalp-components of SER in uncomplicated, unshunted case, 16:157-160
- Centenarians**, and observations of the EEG in 7 cases, 17:146-151
- Central localization** and, mid-line epileptiform activity in adults, 16:54-59
  - midline spikes, 15:232-236
  - spindle activity in the waking EEG with hemispheric glioblastoma, 17:1-5
  - (See also Parietal)
- Central nervous system** and, Cannon and Rosenbluth's law of denervation, 17:No. 4, VII, VIII
- Cerebellar syndrome** and, cerebral electrical fields, 18:142-146
- Cerebral electrical field mapping** and, cerebellar syndrome, 18:142-146
  - (See also Brain electrical activity mapping)
- Cerebral fat embolism**, and periodic EEG patterns, 14:27-34
- Cerebral function monitor (CFM)**, and neonates, 15:185-192
- Cerebral palsy** and, neonatal meningitis, 14:67-77
  - spikes, midline, 15:232-236
- Cerebrovascular disease**, (See Vascular disease)
- Chemotherapy**, and acute lymphocytic leukemia 16:208-212
- CHEERS**, and cerebral fat embolism, 14:27-34
- Children** and, acute lymphoblastic leukemia myelin disturbance, VEP, 18:68-73
  - acute lymphocytic leukemia with fatal subacute immunosuppressive measles encephalitis, 14:214-220
  - acute lymphocytic leukemia and sequential



- quantitative EEG analysis, 16:208-212
- benign focal epilepsy, individual and intrafamilial multifocality of spikes, 18:169-172
- beta activity development, EEG-topographical analysis, 18:120-123
- coma, 14:207-213
- cyclic variation of sleep spindles in mentally retarded, 15:40-46
- EP as outcome predictors in drowning, 16:183-191
- head injury, 18:10-11
- integrated delta and spindle components during nocturnal sleep in the mentally retarded, 17:24-29
- learning disabilities, EEG characteristics, 18:93-101
- Lennox-Gastaut syndrome, frontiers, 17:117-126
- management for EEG, 15:202-207
- myelin disturbance with acute lymphoblastic leukemia, latency of VEP, 18:68-73
- occipital evoked spike epilepsy, 14:221-224
- parietal epilepsy with evoked and spontaneous spikes, report on siblings, 17:159-161
- reactive burst suppression following asphyxia in infants, 14:138-141
- renal disease, quantitative EEG before and after haemodialysis, 17:195-202
- respiratory pauses and apnea during sleep in mentally retarded, 14:90-95
- twenty-four hour ambulatory EEG monitoring, 15:164-166
- unusual EEG patterns, 15:92-96
- valproic acid treatment of epilepsy, 17:14-20
- (See also Neonates)
- Chloral hydrate** and, EEG management of infants and children, 15:202-207
- Chlorpromazine** and, schizophrenia, treatment resistant, computerized EEG profiles, 18:124-135
- Cholinergic mechanisms** and, epilepsy with auditory event-related potentials, 17:10-13
- Choreoathetosis**, paroxysmal dystonic, and photic induced-driven PLEDs, 15:151-154
- Cirrhosis**, liver, and spindle coma in benzodiazepine toxicity, 16:213-218
- Clobazam** and, absence status, 16:111-118
  - paralysis, ictal, status epilepticus treatment, 18:12-14
- Clonazepam** and, familial photosensitive epilepsy 15:47-52
  - Lennox-Gastaut syndrome with status epilepticus, 18:89-92
  - sleep and waking disturbances in epilepsy, 18:136-141
- Clozapine**, and schizophrenia, treatment resistant, computerized EEG profiles, 18:124-135
- Cognitive dysfunction** and, Alzheimer disease, 15:32-39; 16:48-53
  - ECT, 17:66-77
  - EEG changes in the elderly, 14:78-81
  - electrophysiologic validation of cortical and subcortical dementias, 18:26-33
  - FIRDA, 16:1-10
  - Lafora disease with BAEP, 16:202-207
  - lorazepam and methylphenidate effect on event-related potentials and power spectral analysis in normal subjects, 17:176-180
  - mid-line epileptiform activity in adult EEGs, 16:61-64
  - partial epilepsy, interictal, 17:10-13
  - spectral analysis of interictal EEG in epilepsy, 16:94-97
- Coma** and, alpha rhythm, 15:167-172
  - anoxic encephalopathy with coexistence of ectopic rhythms and periodic EEG pattern, 17:187-194
  - brainstem death, 16:63-71
  - burst suppression pattern, extreme stereotypy, 17:162-168
  - clinical and EP studies in alpha coma, 18:103-113
  - coexistence of alpha, theta and spindle pattern coma, 18:74-80
  - delta status in newborn, 15:125-131
  - EEG in prognosis, 14:47-52
  - electrocerebral silence and nasopharyngeal EEG, 18:233-234
  - EMG artifact attenuation with electrocerebral inactivity, 16:72-76
  - near-drowning with hypsarhythmia, 18:180-186
  - pediatric cases, EEG findings, 14:207-213
  - periodic EEG activity, 15:32-39
  - spindles in benzodiazepine toxicity, 16:213-218
  - spindles in viral encephalitis, 18:34-37
  - subacute spongiform encephalopathy with periodic paroxysmal activities, 18:147-158
  - suppression-burst pattern, epileptiform discharges and alpha-theta pattern, 17:82-88
  - theta pattern, 15:116-121
  - theta pattern in younger adults, 18:54-60
- Complex partial seizures** and, small sharp spikes, 18:114-119
  - sphenoidal EEG recording, 17:139-141
  - status epilepticus, two case reports, 14:225-232
- Computerized EEG profiles** and, schizophrenia, treatment resistant, brain surface activity mapping, 18:124-135
  - (See also Brain electrical activity mapping)
- Computerized EEG scanning (CEEGs)**, See Brain electrical activity mapping
- Computerized tomography (CT)** and, cerebral infarction, EEG comparative study, 18:20-25
  - EEG in neurological diagnosis, 14:116-129
  - fatal subacute immunosuppressive measles

- encephalitis in children with acute lymphocytic leukemia, 14:214-220
- near-drowning, with hypsarrhythmia, clinical, EEG, CT and EP studies, 18:180-186
- neonatal meningitis, 14:67-77
- Conceptual Level Analogy Test**, and tardive dyskinesia, 14:186-201
- Concussion** and, electrophysiological tests, 18:165-167
  - (See also, Trauma, head)
- Continuous high amplitude EEG rhythmic synchronous slowing** (CHERS), and cerebral fat embolism, 14:27-34
- Contraceptives, oral**, and suicide risk potentiation in patients with EEG dysrhythmias, 16:88-93
  - letters to the editor, 18:No. 2, XIII
- Convulsions**, (See Grand mal and Seizures)
- Corpus callosum, agenesis**, and, nocturnal sleep in infants, 17:92-104
  - spectral array of delta and spindle components during sleep in infants, 17:152-157
- Corticoid treatment**, and West's syndrome, 16:122-127
  - (See also ACTH)
- Creutzfeldt-Jacob disease** and, clinical evolution and EEG, 18:211-217
  - EEG with CAT scan, 14:116-129
  - periodic EEG activity, 15:32-39; 18:147-158
- Cutaneous femoris lateralis nerve** and, SEP stimulation in meralgia paresthetica, 14:161-163; 15:No. 1, VI, VIII
- Cyst**, and EEG topography in porencephaly and arachnoid cyst, 16:21-29
- Deafness**, (See Hearing loss)
- Death** and, anoxic encephalopathy with ectopic rhythms and periodic EEG pattern, 17:187-194
  - brain death editorial, tread lightly, treat vigorously, 15:No. 1, V, VI
  - Cannon and Rosenbluth's law of denervation, 17:No. 4, VII
  - diagnosis of brain, brainstem, or cerebral death, 16:63-71
  - electrocerebral silence and nasopharyngeal EEG, 18:233-234
  - failure of Pavulon to provide EMG attenuation for electrocerebral inactivity, 16:72-76
  - suppression-burst pattern with spikes and alpha-theta pattern coma, 17:82-88
  - theta pattern coma in younger adults, 18:54-60
  - (See also Electrocerebral inactivity)
- Degenerative diseases** and, Alzheimer's disease, 15:32-39; 16:48-53; 18:26-33, 164
  - Creutzfeldt-Jakob disease, 14:82-85, 116-129; 15:32-39; 18:147-158, 211-217
  - EEG vs. CT in neurological diagnosis, 14:116-129
  - Huntington's chorea, 17:30-35; 18:26-33
  - parkinsonian patients on levodopa, 14:82-85
  - periodic EEG abnormality, 15:32-39
  - Rett's syndrome with Lennox-Gastaut pattern, 18:187-190
  - Wilson's disease, with liver transplantation, 16:192-194
- Delirium**, and FIRDA, 16:1-10
- Delirium tremens** and, EEG pattern, 16:30-32
  - spindle coma in benzodiazepine toxicity, 16:213-218
- Delta activity** and, Alzheimer's disease, 18:164
  - anterior bradyrhythmia compared to FIRDA, 16:1-10
  - centenarians, 17:146-151
  - cerebellar syndrome, cerebral electrical field mapping, 18:142-146
  - cerebral fat embolism, 14:27-34
  - cerebral infarction, 18:20-25
  - children with end stage renal disease before and after haemodialysis, 17:195-202
  - coma, (See Coma)
  - congenital hydrocephalus, 15:22-31
  - ECT, analysis of frontotemporal and nasopharyngeal spectral energy, 17:66-77
  - ECT with spectral energy distributions and methohexital, Innovar and ketamine anesthetics, 17:203-215
  - EEG vs. CT in neurological diagnosis, 14:116-129
  - EEG topography in porencephaly and arachnoid cyst, 16:21-29
  - fever in adult patients, 18:85-88
  - FIRDA, (See Frontal intermittent rhythmic delta activity)
  - glioblastoma, with spindle activity awake, 17:1-5
  - lithium specific EEG changes, 18:46-53
  - nocturnal sleep in mentally retarded children, 17:24-29
  - pediatric cases and coma, 14:207-213
  - periodic abnormalities, 15:32-39
  - pial-dural arteriovenous malformation, FIRDA, 18:227-232
  - post-traumatic syndrome, 15:214-221
  - schizophrenia, treatment resistant, brain surface activity mapping, 18:124-135
  - sleep in infants with congenital cerebral malformation, compressed spectral array, 17:152-157
  - tardive dyskinesia, 17:30-35
  - topographic mapping, artifacts, 18:61-67
  - subclinical status in newborn, an unfavorable prognostic sign, 15:125-131
  - (See also Slow activity)
- Dementia** and, Alzheimer disease, 15:32-39; 16:48-53
  - cortical and subcortical, electrophysiologic validation, 18:26-33
  - EEG changes in the elderly, 14:78-81



- FIRDA, 16:1-10
- Lafora disease, with BAEP, 16:202-207
- mid-line epileptiform activity in adult EEGs, 16:54-61
- periodic EEG activity, 15:32-39
- subacute spongiform encephalopathy with periodic paroxysmal activities, 18:147-158
- Demyelinating disorder** and, hypsarhythmia with ACTH treatment, 16:178-182
- Denervation** and, Cannon and Rosenblueth's law, editorial, 17:No. 4, VII, VIII
- Depressed amplitude** and, congenital hydrocephalus, 15:22-31
- diagnosis of neurological disorders, 14:116-129
- neonatal meningitis, 14:67-77
- (See also Asymmetry, Electroencephalographic inactivity and Low voltage)
- Depression** and, lithium-specific EEG changes, 18:46-53
- REM latency and distortion of sleep patterns, 18:167
- suicide risk in patients with dysrhythmias taking oral contraceptives, 16:88-93
- Depth recording** and, Lennox-Gastaut syndrome, 18:191-200
- seizure control by mesothalamic reticular stimulation, 17:52-60
- seizures with pain, 16:195-201
- seizure surgery, 15:83-91
- Developmental quotient** and, infants with congenital cerebral malformation, 17:92-104
- Diabetes**, and bursts of rhythmic temporal theta, 15:145-150
- Diazepam** and, hypnogenic paroxysmal dystonia, 16:136-142
- seizure management in mentally retarded epileptics, 14:164-172
- spindle coma with toxicity, 16:213-218
- subclinical delta status in newborn, 15:125-131
- Dilantin (diphenylhydantoin)** and, absence status termination and suppression of interictal bursts, 17:173-175
- ECT modification, 17:203-215
- infusion in seizure patients, 15:173-177
- interictal EEG prediction of response, 16:131-135
- mentally retarded epileptics, minimal medication, 14:164-172
- neonatal status epilepticus, 15:197-201
- sleep and waking disturbances in epilepsy, 18:136-141
- valproic acid, effect on serum levels, 16:104-110
- Dipropyl acetic acid**, (See Valproic acid)
- Dizziness** and, BAEP, 15:110-115
- electrophysiological tests and post-concussion syndromes, 18:165-167
- post-traumatic syndrome, 15:214-221
- Dopamine agonist**, and chronic hepatic encephalopathy, effect on EEG, 14:86-89
- tardive dyskinesia, 17:30-35
- Drowning**, and hypsarhythmia-infantile spasms, clinical, EEG, CT and EP studies, 18:180-186
- survival in young children after drowning, brain evoked potentials as outcome predictors, 16:183-191
- Drowsiness** and, episodic anterior drowsy theta in adults, 17:135-138
- low voltage EEG, 14:102-105
- normal patterns in children, 15:92-96
- Drugs** and, absence status, 16:111-118
- alcoholism, 14:8-19; 15:102-109; 16:30-32
- anesthetic effect on EEG, (See Anesthesia)
- antiepileptic, (See Antiepileptic medication)
- beta activities development in children, 18:120-123
- coma, pediatric cases, 14:207-213
- delta status in newborn, 15:125-131
- drowning survival in children, 16:183-191
- EMG attenuation, 16:72-76
- familial photosensitive epilepsy, 15:47-52
- fast activity, 15:237-241
- hepatic encephalopathy, 14:86-89
- hypnagogic paroxysmal dystonia, 16:136-142
- hypsarhythmia, 15:140-144; 16:120-127, 178-182
- interictal EEG, 16:131-135
- intoxication, 14:207-213; 15:32-39, 102-109, 173-177; 16:213-218
- Lennox-Gastaut syndrome, 17:117-126
- Lennox-Gastaut syndrome with status epilepticus, 18:89-92
- lithium-specific pathological EEG changes, replication of earlier investigative results, 18:46-53
- lorazepam and methylphenidate effect on event-related potentials and spectral analysis in normal subjects, 17:176-180
- low voltage EEG, 14:102-105
- lymphocytic leukemia, 16:208-212
- management of infants and children, 15:202-207
- neonatal seizures, 15:125-131, 133-139, 197-201
- nitrous oxide abuse, 16:143-148
- parkinsonian patients on levodopa, 14:82-85
- phenytoin infusion in seizure patients, 15:173-177
- phenytoin termination of absence status, 17:173-175
- schizophrenia, treatment resistant, mapping, 18:124-135
- seizure management in institutionalized mentally retarded, 14:164-172
- sleep and waking disturbances in epilepsy, 18:136-141
- spindle coma in benzodiazepine toxicity,

- 16:213-218
- stuttering and seizures in adults, 17:142-145
  - tardive dyskinesia, 15:222-231, 17:30-35
  - valproic acid effect on Dilantin serum levels, 16:104-110
  - valproic acid treatment of epilepsy in children, 17:14-20
  - writing epilepsy, 17:21-23
- Dysfluency**, and, stuttering and seizures, adult onset, 17:142-145
- Dyskinesia, tardive** (see Tardive dyskinesia)
- Dystonia, hypnogenic paroxysmal**, and, epilepsy or sleep disorder, case report, 16:136-142
- Editorials**, brain death, tread lightly, treat vigorously, 15:No. 1, V-VI
- epilepsy and driving, 14:No.3, V-VI
  - historical notes, EEG interpretation and technique, 14:No. 2, V, No. 3, VII-VIII
  - international standards for EEG instruments, 15:No. 3, VI
  - neurovascular update, 18:No. 2, IX-XI
  - Richard Jung in memoriam, 18:No. 2, VII-VIII
  - topographic mapping, 18:No. 2, IX
- Electrocerebral inactivity** and, brain death, 15:No.1,V
- coma after trauma, 15:116-121
  - failure of Pavulon to provide EMG attenuation, 16:72-76
  - following delta status in newborn, 15:125-131
  - nasopharyngeal EEG, 18:233-234
  - pediatric cases of coma, 14:207-213
  - prognosis of coma, 14:47-52
- Electroconvulsive therapy (ECT)** and, III, analysis of frontotemporal and nasopharyngeal spectral energy, 17:66-77
- IV, spectral energy distributions with methohexital, Innovar and ketamine anesthetics, 17:203-215
- Electrocorticography** and, information derived from intraoperative recordings during seizure surgery, 15:83-91
- Lennox-Gastaut syndrome, 18:191-200
  - seizure control by mesothalamic reticular stimulation, 17:52-60
  - seizures with pain, 16:195-201
- Electrode placement** and, APEEGE system, 15:1-21
- electrocerebral silence and nasopharyngeal EEG, 18:233-234
  - historical notes, 14:No. 2, V, No.3, VII, VIII
  - Neo-EEG system, 15:71-82
- Electronystagmography** and, electrophysiological tests and post-concussion syndrome, 18:165-167
- Embolism**, cerebral fat, and periodic EEG patterns, 14:27-34
- EMG** and, failure of Pavulon to attenuate, with electrocerebral inactivity, 16:72-76
- Encephalitis** and, abnormal EEG patterns in children, 15:92-96
- EEG vs. CT in diagnosis, 14:116-129
  - herpes simplex virus, with periodic sharp complexes, survival in an elderly man, 18:2-9
  - pediatric cases of coma, 14:207-213
  - periodic EEG abnormality, 14:116-129; 15:32-39
  - spindle coma, 18:34-37
  - subacute immunosuppressive measles encephalitis and acute lymphocytic leukemia, 14:214-220
- Endarterectomy**, carotid, and assessment of variability of early scalp-components of the SER in uncomplicated, unshunted case, 16:157-160
- Epilepsia partialis continua** and, electrocorticography, 15:83-91
- pertussis infection, 16:45-47
- Epilepsy**, (See Seizures)
- Epileptiform discharges**, and success of EEG in confirming epilepsy, 16:98-103
- suppression-burst pattern with alpha-theta pattern coma, 17:82-88
  - (See also Spikes and Spike and Wave)
- Episodic anterior drowsy theta (EADT)**, and, adults, 17:135-138
- Ethosuccinimide** and, termination of absence status, 17:173-175
- Evoked potentials** and acupuncture, 14:53-56
- alcohol abuse, pattern reversal VEP, effects of age, 15:102-109
  - alcoholics, short latency, 14:8-19
  - alpha coma, 18:103-113
  - Alzheimer disease, event related, 16:48-53
  - antisocial personality, 16:91-93
  - brain-injured patients with rage attacks and self-injurious behavior, 17:169-172
  - brainstem death, 16:63-71
  - carotid endarterectomy, uncomplicated, unshunted, 16:157-160
  - cerebellar syndrome, cerebral electrical field mapping, 18:142-146
  - dementias, cortical and subcortical, 18:26-33
  - dizziness, 15:110-115
  - drowning outcome predictors in young children, 16:183-191
  - drowning with hypsarrhythmia-infantile spasms, 18:180-186
  - electrophysiological tests and post-concussion syndrome, 18:165-167
  - epilepsy, 16:33-38; 17:10-13
  - false negative intraoperative SEP with simultaneous bilateral stimulation, 17:6-9
  - FIRDA, 14:135-137
  - hepatic injury status, 18:15-19
  - hydrocephalus, congenital, 15:22-31
  - Lafora disease, 16:202-207
  - liver transplantation, monitoring cerebral

- function, 16:192-194
- long-latency auditory event-related potentials in epilepsy, 17:10-13
- lumbar disease, 14:152-160
- meralgia paresthetica, 14:161-163; 15: No.1,VII
- modified auditory brainstem responses (MABR), Part I - rationale and normative study, 18:218-226
- monocular VEP, crossed asymmetry, 16:77-82
- myelin disturbance in children treated for acute lymphoblastic leukemia, latency of VEP, 18:68-73
- neonatal meningitis, 14:67-77
- nitrous oxide abuse, 16:143-148
- normal subjects, effect of lorazepam and methylphenidate on event-related potentials and spectral analysis, 17:176-180
- operating room, book review, 18:No. 2, XII
- pattern reversal VEP, latency changes with gender and age, 18:159-162
- psychiatric patients, pattern reversal, 15:178-183
- renal hemodialysis, 14:202-206, 15:97-101
- sensory-neural hearing loss, 15:159-163
- spinal cord lesions, 14:142-151, 173-185
- status epilepticus, non-convulsive, 16:33-38
- surgical monitoring, 16:83-87
- thoracic outlet syndrome, 17:112-116
- tibial, peroneal, sural nerve stimulation, 16:149-156
- trigeminal nerve stimulation, 14:61-66
- Evoked spike epilepsy** and, benign infantile epilepsies, 16:39-44
  - law of denervation, 17:No.4, VII
  - occipital, in childhood, 14:221-224
  - parietal epilepsy, report on siblings with possible genetic transmission, 17:159-161
  - sharp wave excitability, 17:127-134
- Extrapyramidal signs** and, clinical EEG variables suggesting side effect risk, 18:173-179
- Eye-closure**, and absence status, 16:111-118
- Eye disorder**, (see Visual disorder)
- Familial disorders** and, benign focal epilepsy of childhood, individual and intrafamilial multifocality of spikes, 18:169-172
  - benign infantile epilepsies with evoked spikes, 16:39-44
  - grand mal discharge, 15:155-158
  - mu rhythm, 17:44-50
  - parietal epilepsy with evoked and spontaneous spikes, report on siblings with possible genetic transmission, 17:159-161
  - paroxysmal dystonic choreoathetosis, with PLEDs, 15:151-154
  - photosensitive epilepsy, effectiveness of clonazepam, 15:47-52
  - Rett's syndrome with Lennox-Gastaut pattern, 18:187-190
  - tuberous sclerosis, long-term follow-up, 16:219-224
- Fast activity** (see Beta activity)
- Fat embolism**, cerebral, and periodic EEG patterns, 14:27-34
- Febrile convulsions**, and benign infantile epilepsies with evoked spikes, 16:39-44
  - epileptic syndromes in childhood, book review, 17: No. 3, VII
- Fever** and, EEG in adult patients, 18:85-88
- Fibrositis syndrome**, and, somatic complaints with disturbed sleep, 18:81-84
- FIRDA**, (See Frontal intermittent rhythmic delta activity)
- Focal seizures** and, *epilepsia partialis continua* following pertussis infection, 16:45-47
  - multiple sclerosis with PLEDs, 18:38-40
  - natural history of hypsarhythmia, 16:128-130
  - paralysis, ictal, EEG confirmation of its epileptic nature, 18:12-14
  - stuttering, adult onset, 17:142-145
  - subacute immunosuppressive measles encephalitis with acute lymphocytic leukemia, 14:214-220
  - subacute spongiform encephalopathy, 18:147-158
- Fourteen and six per second positive spikes** and, controversial patterns in children, 15:92-96
  - head injury in childhood, 18:10-11
  - post-traumatic syndrome, 15:214-221
- Frontal intermittent rhythmic delta activity** and, anterior bradyrhythmia, significance of the two patterns, 16:1-10
  - cerebral fat embolism, 14:27-34
  - cerebral infarction, comparative EEG and CT scan study, 18:20-25
  - clinical significance, 14:135-137
  - periodic EEG abnormalities, correlations, 15:32-39
  - pial-dural arteriovenous malformation, 18:227-232
- Frontal localization** and, anterior bradyrhythmia, 16:1-10
  - centenarians, slowing, 17:146-151
  - cerebral fat embolism, slowing, 14:27-34
  - complex partial status epilepticus, spikes, 14:225-232
  - depth EEG studies in Lennox-Gastaut syndrome, 18:191-200
  - ECT, analysis of spectral energy, 17:66-77
  - ECT with methohexital, Innovar and ketamine anesthetics, slowing, 17:203-215
  - electrocorticograms, spikes, 15:83-91
  - electrode placement systems, 15:1-21, 71-82
  - FIRDA, 14:27-34, 135-137; 15:32-39; 16:1-10; 18:20-25, 227-232
  - glioma of the sylvian region with post-ictal gustatory hallucination, microspikes, 17:89-91

- sleep humps in hydrocephalus, 15:22-31
- Fuster, Bartolome**, notice of death, accomplishments, 14:No.3,V
- Generalized periodic slow wave complexes (GPSC)**, correlations of periodic EEG activity, 15:32-39
- Genetic transmission** and, benign focal epilepsy of childhood, individual and intrafamilial multifocality of spikes, 18:169-172
  - parietal epilepsy with evoked and spontaneous spikes in siblings, 17:159-161
  - (See also Familial disorders)
- Geriatric EEG**, and age-dependent incidence of POSTS, 16:11-15
  - photic driving, 14:78-81
  - (See also Adult EEG)
- Gibbs, Erna Leonhardt**, obituary notice, 18:No. 4, VIII
- Glioblastoma**, and, spindle activity in the waking EEG, report of a case with hemispheric glioblastoma, 17:1-5
- Grand mal discharge**, 15:155-158
- Grand mal seizures** and, benign focal epilepsy of childhood, 18:169-172
  - ECT with methohexital, Innovar and ketamine anesthetics, 17:203-215
  - evoked sharp wave excitability, 17:127-134
  - familial photosensitive epilepsy, 15:47-52
  - grand mal discharge, 15:155-158
  - hypsarhythmia, natural history, 16:128-130
  - Lennox-Gastaut syndrome, 17:117-126
  - mid-line spikes, 15:232-236
  - neonatal seizures, 15:133-139, 193-196
  - sleep and waking disturbances in epilepsy, 18:136-141
  - small sharp spikes, 18:114-119
  - West's syndrome, 16:122-127
  - (See also Seizures)
- Guillain Barre syndrome**, and nitrous oxide abuse, with VEP, 16:143-148
- Gustatory hallucinations** and, glioma of the sylvian region with microspikes, 17:89-91
- Hallucinations** and, glioma of the sylvian region, post-ictal gustatory, 17:89-91
- Haloperidol** and, lithium-specific EEG changes, 18:46-53
  - schizophrenia, treatment resistant, computerized EEG profiles, 18:124-135
- Headache** and, absence status control by mesothalamic reticular stimulation, 17:52-60
  - benign focal epilepsy of childhood, 18:169-172
  - electrophysiological tests and post-concussion syndromes, 18:165-167
  - hyperventilation responses in the adult EEG, 17:61-65
  - post-traumatic syndrome, 15:214-221
  - rhythmical temporal theta, 15:145-150
  - stuttering and seizures, adult onset, 17:142-145
- whiplash injury, EEG reappraisal, 18:41-45
- Head Injury** and, childhood, EEG study, 18:10-11
  - (See also Trauma, head)
- Hearing loss** and, auditory EP in surgical monitoring, 16:83-87
  - differentiation between brainstem lesions and sensori-neural hearing loss, with ABER, 15:159-163
- Hemodialysis** and, children with end stage renal disease, quantitative EEG before and after treatment, 17:195-202
  - chronic renal failure, changes in short-latency SEP, 14:202-206
  - pattern reversal EP, 15:97-101
- Hepatic encephalopathy** and, bromocriptin therapy effect on EEG, 14:86-89
  - EEG vs. CT in neurological diagnosis, 14:116-129
  - event related potentials, relationship to injury status, 18:15-19
  - SEP monitoring cerebral functions following liver transplantation, 16:192-194
  - spindle coma in benzodiazepine toxicity, 16:213-218
- Hereditary disorder**, (See Familial disorder)
- Herpes simplex encephalitis**, and EEG periodicity in diagnosis, 14:116-129
  - periodic sharp complexes, survival in an elderly man, 18:2-9
  - spindle coma, 18:34-37
- Hippocampus** and, ischemia with minor slow and sharp activity, 18:201-210
- Holoprosencephaly** and, nocturnal sleep in infants, 17:92-104, 152-157
- Humps, sleep**, and asynchrony with hydrocephalus, 15:22-31
- Huntington's chorea** and, electrophysiological evaluation of cortical and subcortical dementias, 18:26-33
  - tardive dyskinesia, 17:30-35
- Hydranencephaly**, and electrophysiological evaluation of cortical and subcortical dementias, 18:26-33
  - nocturnal sleep in infants, 17:92-104, 152-157
- Hydrocephalus** and, EEG vs. CT in neurological diagnosis, 14:116-129
  - electrophysiologic validation of cortical and subcortical dementias, 18:26-33
  - newborns, neurophysiological evaluation, 15:22-31
  - nocturnal sleep in infants, 17:92-104, 152-157
- Hypertensive encephalopathy**, and, EEG and anatomo-clinical evidences of posterior cerebral damage, 15:53-60
  - FIRDA in pial-dural arteriovenous malformation, 18:227-232

- Hyperventilation** and, EEG characteristics of children with learning disabilities, 18:93-101
- paroxysmal responses in the adult EEG, 17:61-65
- Hypnagogic hypersynchrony**, and normal drowsy patterns in children, 15:92-96
- Hypnogenic paroxysmal dystonia** and, epilepsy or sleep disorder, case report, 16:136-142
- Hypoesthesia**, and, thoracic outlet syndrome, diagnostic importance of SEP, 17:112-116
- Hypoxia** and, drowning survival in children, with EP, 16:183-191
- ectopic rhythms and periodic EEG pattern, 17:187-194
  - neonatal seizures, 14:67-77; 15:133-139, 193-196
  - reactive burst suppression in infants, 14:138-141
  - suppression-burst pattern, epileptiform discharges, and alpha-theta pattern coma, 17:82-88
  - theta pattern coma in younger adults, 18:54-60
  - (See also Apnea)
- Hypsarhythmia** and, ACTH treatment of infantile spasms, Sorel's early success and ongoing benefits, 16:120-121
- diminishing incidence, 16:178-182
  - infection as a possible etiology, 17:78-81
  - Lennox-Gastaut syndrome, 15:61-68, 17:117-126
  - natural history, 16:128-130
  - near-drowning, clinical, EEG, CT and EP studies, 18:180-186
  - pyretotherapy as treatment, 15:140-144
  - treatment of West's syndrome, 16:122-127
  - tuberous sclerosis, 16:219-224
  - valproic acid treatment, 17:14-20
- Infantile spasms**, (See Hypsarhythmia)
- Infants**, (See Neonates and infants)
- Infarction, cerebral** and, EEG and CT scan comparative study, 18:20-25
- (See also Vascular disease)
- Infectious disease** and, EEG vs. CT in neurological diagnosis, 14:116-129
- EEG patterns in children, 15:92-96
  - fever in adult patients, EEG findings, 18:85-88
  - herpes simplex virus encephalitis with periodic sharp complexes, survival in an elderly man, 18:2-9
  - hypsarhythmia, diminishing incidence, 16:178-182
  - hypsarhythmia, possible etiology, 17:78-81
  - pediatric cases of coma, 14:207-213
  - periodic EEG abnormality, 14:116-129; 15:32-39
  - pertussis with epilepsy partialis continua, 16:45-47
  - prognostic value of EEG in neonates, 14:67-77
- spindle coma in viral encephalitis, 18:34-37
  - subacute immunosuppressive measles encephalitis and acute lymphocytic leukemia, 14:214-220
- Innovar** and, ECT with EEG spectral energy distributions, 17:203-215
- Instrument standards**, EEG, 15:No. 3, VI
- Interleukin-1** and, EEG in adult patients with fever, 18:85-88
- International 10-20 System**, and APEEGE system of derivation, 15:1-21
- Irritative foci** and, Cannon and Rosenblueth's law of denervation, 17:No. 4, VII, VIII
- (See also Spikes)
- Isoelectric EEG**, (See Electroencephalographic inactivity)
- Jacksonian seizures** and, Cannon and Rosenblueth's law of denervation, 17:No. 4, VIII
- Jung, Richard**, in memoriam, 18:No. 2, VII-VIII
- K-complex**, and seizures with pain, 16:195-201
- sleep-waking cycle disturbances in vascular brain stem lesions, 14:35-36
- Ketamine (Ketalar)** and, ECT with EEG spectral energy distributions, 17:203-215
- Kidney disease**, (See Renal disease)
- Lacunar state** and, electrophysiologic validation of cortical and subcortical dementias, 18:26-33
- Lafora disease** and, brainstem auditory evoked responses, 16:202-207
- occipital seizures, case documented by EEG, 16:167-170
- Learning-disability** and, EEG characteristics of children, 18:93-101
- Lennox-Gastaut syndrome** and, depth EEG studies, 18:191-200
- frontiers, review 17:117-126
  - grand mal discharge, 15:155-158
  - natural history of hypsarhythmia, 16:128-130
  - Rett's syndrome, 18:187-190
  - status epilepticus, 18:89-92
  - valproic acid treatment, 17:14-20
  - West's syndrome evolution, 15:61-68
- Letters to the Editor**, lymphoblastic leukemia, myelin disturbance in children, with VEP, 18:No. 3, VI, VII
- meralgia paresthetica, SEP, 15:No., VII
  - monopolar vs. bipolar recording, 14:No. 3, VII-VIII
  - oral contraceptives, EEG dysrhythmias and suicide risk, 18:No. 2, XIII
  - theta pattern coma, 16:No. 3, VIII
- Leukemia** and, fatal subacute immunosuppressive measles encephalitis in children, with acute lymphocytic leukemia, clinical EEG and CT features, 14:214-220
- myelin disturbance in children treated for acute lymphoblastic leukemia, latency of



- VEP, 18:68-73; No. 3, VI-VII
- sequential quantitative analysis in children, with acute lymphocytic leukemia, 16:208-212
- Levo-dopa** and, parkinsonian patients, quantitative study of EEG response, 14:82-85
- Librium** and, alcoholics with SEP, 14:8-16
- Lightning** and, artifact in the EEG, 17:105-109
- Lithium therapy** and, pathological EEG changes, replication of earlier results, 18:46-53
- tardive dyskinesia, 15:222-225
- Liver disease**, (See Hepatic encephalopathy)
- Liver transplantation** and, SEP monitoring cerebral functions, 16:192-194
- Locked-in syndrome** and, brainstem infarction with sleep disturbances, 14:35-46
- coma with EEG, 14:47-52
- Lorazepam** and, event-related potentials and spectral analysis in normal subjects, 17:176-180
- spindle coma in toxicity, 16:213-218
- Low voltage EEG** and, delirium tremens, 16:30-32
- diagnosis of neurological disorders, 14:116-129
  - hydrocephalus, 15:22-31
  - neonatal meningitis, 14:67-77
  - nocturnal sleep in infants with congenital malformation, 17:92-104
  - overview, 20 years experience, 14:102-105
  - pediatric cases of coma, 14:207-213
  - (See also Electroencephalographic inactivity)
- Lumbar disc disease** and, dermatomally stimulated somatosensory cerebral evoked potentials, 14:152-160
- Lymphocytic leukemia** (See Leukemia)
- Malformations**, brain, and delta and spindle components in compressed spectral array during sleep in infants, 17:152-157
- nocturnal sleep in infants, 17:92-104
- Management**, of infants and children for EEG, 15:202-207
- Mapping**, See Brain electrical activity mapping
- Measles (Rubeola)** and, diminishing incidence of hypsarrhythmia, 16:178-182
- hypsarrhythmia, possible etiology, 17:78-81
- Measles encephalitis**, fatal subacute immunosuppressive, in children with acute lymphocytic leukemia, clinical EEG and CT features, 14:214-220
- Median nerve**, diagnostic importance of SEP in thoracic outlet syndrome, 17:112-116
- Medications**, and seizure management in mentally retarded epileptics, 14:164-172
- (See also Antiepileptic medication and Drugs)
- Medicine, changing practice**, guest editorial, 17: No. 3, VIII-XII
- Memory impairment**, (See Cognitive dysfunction)
- Meningitis** and, pediatric cases of coma, 14:207-213
- prognostic value of EEG in neonates, 14:67-77
- Mental retardation** and, ACTH treatment of infantile spasms, 16:120-121
- antiepileptic medication serum levels, 16:104-110
  - cyclic variation of sleep spindles, in children, 15:40-46
  - grand mal discharge, 15:155-158
  - hypsarrhythmia, natural history, 16:128-130
  - Lennox-Gastaut syndrome, 17:117-126; 18:191-200
  - minimal medication in seizure management, 4½ year follow up, 14:164-172
  - neonatal meningitis, 14:67-77
  - nocturnal sleep integrated delta and spindle components, 17:24-29
  - nocturnal sleep in infants with congenital cerebral malformation, 17:92-104; 152-157
  - respiratory pauses and apnea during nocturnal sleep in infants, 14:90-95
  - Rett's syndrome with Lennox-Gastaut pattern, 18:187-190
  - tuberous sclerosis, 16:219-224
  - very fast beta, 15:237-241
  - West's syndrome, 15:61-68, 140-144; 16:122-127
- Meralgia paresthetica**, and SEP from stimulation of cutaneous femoris lateralis nerve, 14:161-163; 15:No.1,VII
- Metabolic disorder** and, FIRDA, 14:135-137
- Lafora disease, 16:167-170, 202-207
  - periodic abnormalities, 15:32-39
  - periodic lateralized epileptiform discharges, possible role in pathogenesis, 17:36-43
- Methohexital**, (Brevital) and ECT with EEG analysis, 17:203-215
- Methotrexate**, and acute lymphocytic leukemia in children, 16:208-212
- Methylphenidate** and, normal subjects, effect on event-related potentials and spectral analysis, 17:176-180
- Microcephaly** and, nocturnal sleep in infants, 17:92-104
- spectral array of delta and spindle components during sleep in infants, 17:152-157
- Microfiche** and, Clinical EEG, 14:No.2,VII
- data reduction for the EEG laboratory, 14:106-114
- Mid-line localization** and, clinical and EEG correlates, spikes, 15:232-236
- epileptiform activity in adult EEGs, 16:54-59
- Mid-temporal localization** and, rhythmic theta, 15:145-150
- very fast beta, 15:237-241
  - (See also Temporal)
- Migraine** and, paroxysmal hyperventilation



- responses in the adult EEG, 17:61-65
- (See also Headache)
- Minimal temporal slow activity**, and bursts of rhythmical temporal theta, 15:145-150
- Mitten pattern** and, clinical EEG variables suggesting extrapyramidal side effect risk, 18:173-179
- tardive dyskinesia, 14:186-201; 15:222-225; 17:30-35
- Monopolar recording** and, APEEGE system, 15:1-21
- historical notes, 14:No.2,V;No.3,VII,VIII
- Neo-EEG system, 15:71-82
- Mu rhythm** and, electrocorticograms, 15:83-91
- familial occurrence, 17:44-50
- mid-line epileptiform activity in adult EEGs, 16:54-59
- normal patterns in children, 15:92-96
- post-traumatic syndrome, 15:214-221
- Multiple sclerosis** and, misdiagnosis in nitrous oxide abuse, 16:143-148
- PLEDs in a patient with definite disease, 18:38-40
- usefulness of EEG, 18:163-164
- Muscle activity, legs (MAL)**, and frequent arousals in narcoleptics and obstructive sleep apnea patients, 17:181-186
- Myelin disturbance** and, acute lymphoblastic leukemia in children, latency of VEP, 18:68-73
- Myoclonic seizures** and, absence status, 16:111-118
- anoxic encephalopathy, coexistence of ectopic rhythms and periodic EEG pattern, 17:187-194
- Creutzfeldt-Jacob disease, 18:211-217
- familial photosensitive epilepsy, 15:47-52
- Lafora disease with BAER, 16:167-170, 202-207
- Lennox-Gastaut syndrome, 17:117-126
- subacute spongiform encephalopathy, 18:147-158
- suppression-burst pattern, spikes and alpha-theta pattern coma, 17:82-88
- valproic acid treatment in juvenile type, 17:14-20
- West's syndrome, 15:61-68, 140-144
- writing epilepsy, 17:21-23
- Myoclonus, nocturnal** and, frequent arousals in narcoleptics and obstructive sleep apnea patients, 17:181-186
- sleep disturbances, 14:130-134
- Nagler, Dr. Benedict**, in memoriam, 15: No. 2,V
- Narcolepsy**, and disturbed nocturnal sleep, 14:130-134
- muscle activity in the legs with frequent arousals, 17:181-186
- Nasopharyngeal electrodes** and, ECT, analysis of spectral energy, 17:66-77
- ECT with methohexital, Innovar and ketamine anesthetics, slowing, 17:203-215
- electrocerebral silence, 18:233-234
- Neo-electroencephalography (N.E.E.G.)**, basis and use of a true monopolar derivation, 15:71-82
- Neonates and infants**, and, cerebral function monitor studies, 15:185-192
- congenital cerebral malformations, nocturnal sleep, 17:92-104, 152-157
- development of beta activities, EEG-topographical analysis, 18:120-123
- hydrocephalus, congenital, neurophysiological evaluation, 15:22-31
- hypsarhythmia (See Hypsarhythmia)
- management for EEG, 15:202-207
- meningitis, prognostic value of EEG, 14:67-77
- reactive burst suppression following asphyxia, 14:138-141
- respiratory pauses and apnea during sleep in mentally retarded, 14:90-95
- seizures, a reappraisal, 15:133-139
- status epilepticus, clinical aspects, 15:193-196
- status epilepticus, EEG aspects, 15:197-201
- subclinical delta status, 15:125-131
- Neoplasm**, (See Tumor)
- Neuro-Behcet syndrome**, and periodic lateralized EEG abnormality, 15:122-124
- Neuroleptic medication** and, clinical EEG variables suggesting extrapyramidal side effect risk, 18:173-179
- lithium-specific pathological EEG changes, 18:46-53
- schizophrenia, treatment resistant, brain surface activity mapping, 18:124-135
- tardive dyskinesia, 14:186-201; 15:222-225, 226-231
- (See also Drugs)
- Neurological diagnosis**, and EEG vs. CT, 14:116-129
- Neuro-otologic procedures**, and ABR surgical monitoring, 16:83-87
- (See also Auditory evoked potentials)
- Neuropsychological correlates**, and EEG in tardive dyskinesia, 5 year follow up, 14:186-201
- Neurovegetative symptoms** and, unusual patterns in children, 15:92-96
- Newborn**, (See Neonates)
- Nitrous oxide**, and abuse, with pattern-reversal visual evoked potential, 16:143-148
- Nocturnal sleep**, and community hospital patient outcome, 16:16-20
- congenital cerebral malformations, 17:92-104, 152-157
- mentally retarded children, 14:90-95, 17:24-29
- narcolepsy, 14:130-134

- pediatrics, 24 hour monitoring, 15:164-166
- post-traumatic syndrome, 15:214-221
- rapid clinical staging, reliability, 16:16-20
- recording in subject's home, 14:96-101
- vascular brainstem lesions, 14:35-46
- (See also Sleep)

**Normal EEG and, dementias, cortical and subcortical, 18:26-33**

- EEG vs. CT in neurological diagnosis, 14:116-129
- EEG confirming epilepsy, 16:98-103
- patterns in children, 15:92-96
- POSTS and geriatric EEG, 16:11-15
- post-traumatic syndrome, 15:214-221
- suicidal behavior, 14:20-26
- universal APEEGE system, anatomical placement of electrodes, 15:1-21

**Normal subjects and, crossed asymmetry in monocular VEP, 16:77-82**

- modified auditory brainstem responses, Part I - rationale and normative study, 18:218-226
- pharmacodynamic properties of lorazepam and methylphenidate on event-related potentials and power spectral analysis, 17:176-180
- SEP stimulation of tibial, peroneal, and sural nerves, 16:149-156

**Obstructive sleep apnea and, muscle activity in the legs with frequent arousals, 17:181-186**

- (See also Apnea)

**Occipital localization and, APEEGE system of electrode placement, 15:1-21**

- benign focal epilepsy of childhood, spikes, 18:169-172
- benign infantile epilepsies, evoked spikes, 16:39-44
- evoked spike epilepsy in childhood, 14:221-224
- hypsarrhythmia, natural history, spikes, 16:128-130
- Lafora disease, spikes, 16:167-170
- law of denervation, spikes, 17:No. 4, VII
- Lennox-Gastaut syndrome, rare variant, spike and wave, 17:117-126
- normal patterns in children, 15:92-96
- parkinsonian patients on levodopa, alpha, 14:82-85
- POSTS in geriatric EEG, 16:11-15

**Occipital seizures and, benign infantile epilepsies with evoked spikes, 16:39-44**

- evoked spike epilepsy of childhood, 14:221-224
- Lafora disease, a further case documented by EEGs, 16:167-170

**Operative monitoring, (See Depth recording and Surgery)**

**PAFA, periodic anterior fast activity, and grave prognosis, 15:32-39**

**Pain and, absence status with control by**

- mesothalamic reticular stimulation, 17:52-60
- post traumatic syndrome, 15:214-221
- seizures, with slow wave discharges, 16:195-201
- thoracic outlet syndrome, diagnostic importance of SEP, 17:112-116
- whiplash injury, EEG reappraisal, 18:41-45
- (See also Headache)

**Pancuronium bromide (Pavulon), and failure to provide EMG attenuation for electrocerebral inactivity, 16:72-76**

**Paralysis, ictal, and, epileptic nature, EEG confirmation, 18:12-14**

- status epilepticus, spikes, 18:12-14

**Parasomnia, and hypnogenic paroxysmal dystonia, 16:136-142**

**Paresthesia and, nitrous oxide abuse with VEP, 16:143-148**

- SEP from cutaneous femoris lateralis nerve in meralgia paresthetica, 14:161-163; 15:No.1, VII

**Parietal epilepsy and, evoked and spontaneous spikes in siblings, possible genetic transmission, 17:159-161**

**Parietal localization and, benign infantile epilepsies, evoked spikes, 16:39-44**

- cerebral infarction, comparative EEG and CT scan study, spikes, 18:20-25
- epileptiform activity in adult EEGs, 16:54-59
- evoked and spontaneous spikes, report on siblings with possible genetic transmission, 17:159-161
- ictal paralysis, status epilepticus, spikes, 18:12-14
- midline spikes, 15:232-236

**Parkinson's disease and, electrophysiological evaluation of cortical and subcortical dementias, 18:26-33**

- quantitative study of EEG response to levodopa treatment, 14:82-85

**Paroxysmal dysrhythmia and, clinical EEG variables suggesting extrapyramidal side effect risk, 18:173-179**

- success of EEG in confirming epilepsy, 16:98-103
- (See also Spike and wave and Spikes)

**Paroxysmal dystonic choreoathetosis, and photic induced-driven PLEDs, 15:151-154**

**Partial Seizures and, benign infantile epilepsies with evoked spikes, 16:39-44**

- BSAEP, 13:154-161
- epilepsia partialis continua, following pertussis infection, 16:45-47
- epilepsia partialis continua with electrocorticography, 15:83-91
- hyperventilation responses in the adult EEG, 17:61-65
- long-latency auditory event-related potentials, 17:10-13
- parietal epilepsy with evoked and spontan-

- eous spikes in siblings, 17:159-161
- psychomotor status epilepticus, 2 cases, 14:225-232
- sleep and waking disturbances in epilepsy, 18:136-141
- small sharp spikes, 15:208-213; 18:114-119
- sphenoidal EEG recording, 17:139-141
- valproic acid treatment in children, 17:14-20

**Pattern reversal visual evoked potentials** and, effects of age and alcohol abuse, 15:102-109

- psychiatric patients, 15:178-183
- renal hemodialysis, 15:97-101
- (See also Visual evoked potentials)

**Pavulon**, and failure to provide EMG attenuation for recording electrocerebral inactivity, 16:72-76

**Pediatric cases** and, coma, EEG findings, 14:207-213

- 24 hour ambulatory EEG monitoring, 15:164-166
- valproic acid treatment of epilepsy, 17:14-20
- (See also Neonates and Children)

**Pentobarbital** and, coma in near-drowning, with hypsarrhythmia-infantile spasms, 18:180-186

**Perinatal injury** and, grand mal discharges, 15:155-158

- neonatal seizures, 15:133-139
- (See also Hypsarrhythmia and Neonates)

**Periodic anterior fast activity (PAFA)** and, periodic EEG activity, 15:32-39

**Periodic EEG patterns** and, cerebral fat embolism, 14:27-34

- cerebral infarction, 18:20-25
- correlations, 15:32-39
- Creutzfeldt-Jacob disease, 18:211-217
- ectopic rhythms in anoxic encephalopathy, 17:187-194
- EEG vs. CT in neurological diagnosis, 14:116-129
- herpes simplex virus encephalitis, 18:2-9
- lymphocytic leukemia, 14:214-220
- metabolic factors, 17:36-43
- multiple sclerosis, 18:38-40
- Neuro-Behcet syndrome, PLEDs, 15:122-124
- paroxysmal dystonic choreoathetosis, 15:151-154
- subacute spongiform encephalopathy, 18:147-158
- (See also Suppression-burst pattern)

**Periodic lateralized epileptiform discharges** (PLEDs) and, acute lymphocytic leukemia with fatal subacute immunosuppressive measles encephalitis, 14:214-220

- cerebral fat embolism, 14:27-34
- cerebral infarction, comparative EEG and CT scan study, 18:20-25
- herpes simplex virus encephalitis, survival in an elderly man, 18:2-9
- metabolic factors, possible role in pathogenesis, 17:36-43
- multiple sclerosis, patient with definite disease, 18:38-40
- Neuro-Behcet syndrome, 15:122-124
- paroxysmal dystonic choreoathetosis, photic induced, 15:151-154
- periodic EEG activity, 15:32-39

**Periodic paroxysmal activity** and, subacute spongiform encephalopathy, clinical evolution and serial EEG findings in 20 cases, 18:147-158

—(See also Periodic EEG patterns)

**Peroneal nerve** and, SEP stimulation, 16:149-156

**Pertussis infection** and, epilepsy partialis continua, 16:45-47

**Petit mal seizures** and, epileptic syndromes in childhood, book review, 17: No. 3, VII

- familial photosensitive epilepsy, 15:47-52
- hypsarrhythmia, natural history, 16:128-130
- Lennox-Gastaut syndrome, 17:117-126
- mentally retarded, seizure management, 14:164-172
- mesothalamic reticular stimulation, 17:52-60
- status appearing on eye-closure, 16:111-118
- status, termination and suppression of ictal bursts with phenytoin, case report, 17:173-175
- valproic acid treatment in children, impulsive, 17:14-20

**Petit mal variant pattern** and, grand mal discharge, 15:155-158

- hypsarrhythmia, natural history, 16:128-130
- Lennox-Gastaut syndrome, 17:117-126
- Lennox-Gastaut syndrome with status epilepticus, clinical characteristics and management, 18:89-92
- tuberous sclerosis, 16:219-224
- West's syndrome evolution, 15:61-68

**Phenobarbital** and, familial photosensitive epilepsy, 15:47-52

- interictal EEG prediction of response, 16:131-135
- neonatal seizures, 15:133-139, 197-201
- seizure management in institutionalized mentally retarded epileptics, 14:164-172
- sleep and waking disturbances in epilepsy, 18:136-141

**Phenytoin** and, absence status termination and suppression of inter-ictal bursts, case report, 17:173-175

- infusion in seizure patients, EEG spectra, 15:173-177
- interictal EEG prediction of response, 16:131-135
- neonatal status epilepticus, 15:197-201
- sleep and waking disturbances in epilepsy, 18:136-141
- (See also Dilantin)

**Photic stimulation** and, evoked sharp wave excitability, 17:127-134

—familial photosensitive epilepsy, with clonazepam, 15:47-52

—Lafora disease, 16:167-170, 202-207

—occipital evoked spike epilepsy in childhood, 14:221-224

—paroxysmal dystonic choreoathetosis, PLEDs, 15:151-154

—post-traumatic syndrome, 15:214-221

—psychogeriatric diagnosis, photic driving, 14:78-81

—subacute spongiform encephalopathy, 18:147-158

**Photosensitive seizures** and, valproic acid treatment in children, 17:14-20

**Pick's dementia** and, electrophysiologic validation of cortical and subcortical dementias, 18:26-33

**PLEDs**, (See Periodic lateralized epileptiform discharges)

**Polygraphic recordings** and, delta and spindle components during sleep in infants with congenital cerebral malformation, 17:152-157

—muscle activity in the legs with frequent arousals in narcolepsy and obstructive sleep apnea, 17:181-186

—narcolepsy with disturbed nocturnal sleep, 14:130-134

—nocturnal sleep in infants with congenital cerebral malformation, 17:92-104

—nocturnal sleep in mentally retarded children, integrated delta and spindle components, 17:24-29

—recording in subject's home, 14:96-101

—respirograms on mentally retarded infants during sleep, 14:90-95

—vascular brain stem lesions with disturbances of sleep-waking cycle, 14:35-46

**Polysomnography** (See Polygraphic recordings)

**Porencephaly**, and EEG topography, 16:21-29

—nocturnal sleep in infants, 17:92-104, 152-157

**Portable cassette recording**, and EEG monitoring of all night sleep, 16:16-20

—(See also Ambulatory EEG)

**Positive occipital sharp transients of sleep (POSTS)**, and age-dependent incidence in geriatric EEG, 16:11-15

**Post-concussion syndromes** and, electrophysiological tests, 18:165-167

**Prematurity** and, cerebral function monitor in neonates, 15:185-192

—neonatal seizures, 15:133-139, 193-196

**Primidone**, and seizure management in mentally retarded epileptics, 14:164-172

**Pseudoseizures** and, ambulatory EEG, community hospital prospective study of patient

outcome, 16:161-166

—saline activation, clinical EEG and neuropsychiatric observations, 16:171-176

—success of EEG in confirming epilepsy, 16:98-103

—24 hour ambulatory EEG in pediatrics, 15:164-166

**Psychiatric disorder** and, complex partial status epilepticus, 14:225-232

—electroconvulsive therapy, 17:203-215

—extrapyramidal side effect risk, clinical EEG variables, 18:173-179

—FIRDA compared to anterior bradyrhythmia, 16:1-10

—lithium-specific EEG changes, replication of earlier results, 18:46-53

—low voltage EEG, 14:102-105

—pattern reversal evoked potentials, 15:178-183

—photic driving in elderly, 14:78-81

—saline activation of pseudoepileptic seizures, 16:171-176

—schizophrenia, treatment resistant, brain surface activity mapping, 18:124-135

—sleep patterns distortion, 18:167

—small sharp spikes, possible relationship to epilepsy, 18:114-119

—success of EEG in confirming epilepsy, 16:98-103

—suicide with EEG, 14:20-26

—suicide risk in patients with dysrhythmias taking oral contraceptives, 16:88-93; 18: No. 2, XIII

—tardive dyskinesia, 14:186-201; 15:222-231; 17:30-35

—very fast beta activity, 15:237-241

**Psychogeriatric diagnosis**, and photic driving, 14:78-81

**Psychometric tests**, and psychiatric patients with tardive dyskinesia, 14:186-201

**Psychomotor seizures** and, small sharp spikes, 18:114-119

—sphenoidal recording, 17:139-141

—status epilepticus, 14:225-232

**Pyrethotherapy**, treatment in West's syndrome, 15:140-144

**Race factor**, and bursts of rhythmical temporal theta, 15:145-150

**Radiation therapy**, and acute lymphocytic leukemia of children, 16:208-212

**Rage attacks** and, brain-injured patients with BAEPs, 17:169-172

**Recording techniques** and, APEEGE system, 15:1-21

—Neo-EEG system, 15:71-82

**Reflex epilepsy** and, writing and typing, 17:21-23

**REM sleep** and, depression, latency changes and distortion of sleep patterns, 18:167

—infants with congenital cerebral malfor-

- mation, 17:92-104
- mentally retarded, 14:90-95
- narcolepsy, 14:130-134
- sleep and waking disturbances in epilepsy, 18:136-141
- sleep studies in the home, 14:96-101
- somatic complaints, 18:81-84
- spindles in mentally retarded children, 15:40-46
- vascular brainstem lesions, 14:36-46
- (See also Sleep)
- Renal disease** and, children with end stage renal disease before and after haemodialysis, quantitative EEG, 17:195-202
- pattern reversal evoked potentials, 15:97-101
- short-latency SEP in chronic renal failure, 14:202-206
- Respiratory pauses** and, apnea during nocturnal sleep in mentally retarded infants, 14:90-95
- (See also Apnea)
- Reticular activating system** and, BAEP in antisocial personality, 16:91-93
- BAEPs in brain-injured patients with rage attacks and self-injurious behavior, 17:169-172
- neurophysiologic mechanisms in ECT seizures, 17:203-215
- seizure control by mesothalamic reticular stimulation, 17:52-60
- seizures with pain, 16:195-201
- Rett's syndrome** and, Lennox-Gastaut pattern, 18:187-190
- Rhythmic mid-temporal activity** and, theta, 15:145-150
- very fast beta, 15:237-241
- (See also PLEDs)
- Rolandic localization**, and benign infantile epilepsies, evoked spikes, 16:39-44
- electrocorticograms, 15:83-91
- evoked and spontaneous spikes in siblings, 17:159-161
- mu rhythm, 17:44-50
- Saline activation** and, pseudoepileptic seizures, clinical EEG and neuropsychiatric observations, 16:171-176
- Schizophrenia** and, computerized EEG profiles of haloperidol, chlorpromazine, clozapine and placebo in treatment resistant cases, 18:124-135
- pattern reversal evoked potentials, 15:178-183
- tardive dyskinesia, 15:222-225, 226-231; 17:30-35
- Sedation**, and management of infants and children for EEG, 15:202-207
- Seizure discharges**, (See Spikes, Spike and wave)
- Seizures** and, absence status appearing on eye-closure, 16:111-118
- absence status control by mesothalamic reticular stimulation, 17:52-60
- ACTH treatment of infantile spasms, 16:120-121
- ambulatory EEG, 15:164-166; 16:161-166
- anoxic encephalopathy, coexistence of ectopic rhythms and periodic EEG pattern, 17:187-194
- antiepileptic drug monotherapy, interictal EEG prediction of response, 16:131-135
- benign focal epilepsy of childhood, individual and intrafamilial multifocality of spikes, 18:169-172
- benign infantile epilepsies with evoked spikes, 16:39-44
- brain-injured patients with rage attacks and self-injurious behavior, 17:169-172
- burst suppression in infants following asphyxia, 14:138-141
- Cannon and Rosenblueth's law of denervation, 17:No. 4, VII, VIII
- cerebrotendinous xanthomatosis, 13:89-96
- complex partial, sphenoidal recording, 17:139-141
- complex partial and small sharp spikes, 18:114-119
- complex partial status epilepticus, 14:225-232
- computerized spectral analysis of interictal EEG, 16:94-97
- delta status in newborn, 15:125-131
- ECT, 17:66-77, 203-215
- EEG vs. CT in diagnosis of neurological disorders, 14:116-129
- electrocorticography, 15:83-91
- epilepsia partialis continua following pertussis, 16:45-47
- epileptic syndromes in childhood, book review, 17: No. 3, VII
- episodic anterior drowsy theta in adults, 17:135-138
- evoked sharp wave, excitability, a case study, 17:127-134
- familial photosensitive epilepsy, effectiveness of clonazepam, 15:47-52
- grand mal discharge, 15:155-158
- gustatory hallucination with glioma of the sylvian region, 17:89-91
- hypertensive encephalopathy, 15:53-60
- hyperventilation responses in the adult EEG, 17:61-65
- hypnogenic paroxysmal dystonia, 16:136-142
- infantile spasms (See Hypsarhythmia)
- Lafora disease, 16:167-170, 202-207
- Lennox-Gastaut syndrome, 15:61-68, 17:117-126; 18:29-92, 191-200
- Lennox-Gastaut syndrome depth EEG studies, 18:191-200
- Lennox-Gastaut syndrome with status epilepticus, 18:89-92



- long-latency auditory event-related potentials, 17:10-13
  - medication, (See Antiepileptic medication)
  - mentally retarded, minimal medications in institutionalized cases, 14:164-172
  - mesothalamic reticular stimulation, 17:52-60
  - mid-line spikes, 15:232-236
  - multiple sclerosis, usefulness of EEG, 18:163-164
  - multiple sclerosis with PLEDs, 18:38-40
  - near-drowning, with hypersarhythmia-infantile spasms, 18:180-186
  - neonates, 15:125-131, 133-139, 193-201
  - neonatal meningitis, 14:67-77
  - neurophysiological basis of epilepsy, 18:163
  - non-convulsive, with EP, 16:33-38
  - occipital evoked, in childhood, 14:221-224
  - pain, with slow discharges, 16:195-201
  - paralysis, ictal, EEG confirmation, 18:12-14
  - parietal, with evoked and spontaneous spikes, report on siblings, 17:159-161
  - paroxysmal dystonic choreoathetosis, 15:151-154
  - pediatric cases of coma, 14:207-213
  - phenytoin infusion, EEG spectra, 15:173-177
  - phenytoin termination of absence status and suppression of interictal bursts, 17:173-175
  - PLEDs, 15:32-39, 151-154; 17:36-43
  - pseudoseizures, (See Pseudoseizures)
  - saline activation, 16:171-176
  - sleep and waking disturbances in epilepsy, 18:136-141
  - small sharp spikes, 15:208-213; 18:114-119
  - sphenoidal EEG recording in complex partial seizures, 17:139-141
  - status epilepticus, (See Status epilepticus)
  - stuttering, adult onset, 17:142-145
  - subacute immunosuppressive measles encephalitis with acute lymphocytic leukemia, 14:214-220
  - subacute spongiform encephalopathy, 18:147-158
  - success of EEG in confirmation, revisited, 16:98-103
  - suppression-burst pattern, spikes and alpha-theta pattern coma, 17:82-88
  - therapeutic drug monitoring, 14:No.3,V,VI
  - treatment, epileptic syndromes in childhood, book review, 17: No. 3, VII
  - tuberous sclerosis, long-term follow-up, 16:219-224
  - valproic acid and diphenylhydantoin, effect on serum levels, 16:104-110
  - valproic acid treatment in children, 17:14-20
  - very fast activity, 15:232-236
  - West's syndrome, (See West's syndrome)
  - whiplash injury, 18:41-45
  - writing epilepsy, 17:21-23
- Senile dementia**, and event-related evoked potentials in Alzheimer disease, 16:48-53
- periodic EEG activity, 15:32-39
  - (See also Dementia)
- SEP**, (See Somatosensory evoked potentials)
- Sex factor** and, bursts of rhythmic temporal theta, 15:145-150
- pattern reversal VEP, latency changes, 18:159-162
  - tardive dyskinesia, 15:226-231
  - unusual EEG patterns in children, 15:92-96
- Sharp waves** and, evoked sharp wave excitability, a case study, 17:127-134
- herpes simplex virus encephalitis, survival in an elderly man, periodic, 18:2-9
  - periodic EEG activity, 15:32-39
  - rhythmic temporal theta, 15:145-150
  - stuttering and seizures, adult onset, 17:142-145
  - temporal minor slow activity with cerebrovascular disorder, 18:201-210
  - (See also Spikes)
- Six per second spike and wave discharges** and, post-traumatic syndrome, 15:214-221
- unusual EEG patterns in children, 15:92-96
  - whiplash injury, 18:41-45
- Sleep** and, alpha coma, clinical and EP studies, 18:103-113
- apnea in mentally retarded infants, 14:90-95
  - Cannon and Rosenbluth's law of denervation, 17:No. 4, VII
  - cerebral function monitor studies in neonates, 15:185-192
  - coma, pediatric cases, 14:207-213
  - congenital hydrocephalus, 15:22-31
  - delta and spindle components in infants with congenital cerebral malformation, 17:152-157
  - depression, distortion of sleep patterns, 18:167
  - EEG findings in adult patients with fever, 18:85-88
  - electrode placement, APEEGE system, 15:1-21
  - epilepsy, disturbances in sleep and waking, 18:136-141
  - epilepsy, EEG in confirmation, 16:98-103
  - FIRDA compared to anterior bradyrhythmia, 16:1-10
  - hypnogenic paroxysmal dystonia, 16:136-142
  - infants with congenital cerebral malformation, 17:92-104, 152-157
  - integrated delta and spindle components in mentally retarded children, 17:24-29
  - Lennox-Gastaut syndrome, 17:117-126
  - mental retardation, 14:90-95, 15:40-46, 17:24-29
  - microspikes with post-ictal gustatory hallucination and glioma of the sylvian region, 17:89-91



- mid-line spikes, 15:232-236
- narcolepsy, 14:130-134; 17:181-186
- neonatal meningitis, cycles, 14:67-77
- obstructive sleep apnea, 17: 181-186
- POSTS in geriatric EEG, 16:11-15
- post-traumatic syndrome, 15:214-221
- recordings in subject's home, 14:96-101
- reliability of rapid clinical staging of all night EEG recordings, 16:16-20
- REM, (See REM sleep)
- seizures with pain, arousal slowing, 16:195-201
- sleep latency testing, a measure of daytime sleepiness, 18:164
- somatic complaints, 18:81-84
- sphenoidal EEG recording in complex partial seizures, 17:139-141
- spindle activity awake and asleep, with glioblastoma, 17:1-5
- spindle coma, (See Spindles, sleep)
- spindle components in mentally retarded children, cyclic variation, 15:40-46
- spindles in viral encephalitis coma, 18:34-37
- subclinical delta status in newborn, 15: 125-131
- vascular brain stem lesions, sleep-waking cycle disturbances, 14:35-46
- Slow activity** and, Alzheimer's disease, 18:164
- anoxic encephalopathy, 17:187-194
- anterior bradyrhythmia compared with FIRDA, 16:1-10
- antiepileptic drug monotherapy, interictal EEG prediction of response, 16:131-135
- asphyxia in infants, 14:138-141
- brainstem death, 16:63-71
- cerebellar syndrome, cerebral electrical field mapping, 18:142-146
- centenarians, 17:146-151
- cerebral fat embolism, 14:27-34
- cerebral infarction, comparative EEG and CT scan study, 18:20-25
- cerebrovascular disorder, temporal minor slow and sharp activity, 18:201-210
- children with end stage renal disease before and after haemodialysis, 17:195-202
- children with learning disabilities, 18:93-101
- coma, 14:47-52, 207-213; 15:32-39, 116-121; 17:82-88, 186-193; 18:34-37, 54-60, 74-80
- complex partial status epilepticus, 14: 225-232
- Creutzfeldt-Jacob disease, periodic sharp waves, 18:211-217
- dementias, cortical and subcortical, 18:26-33
- ECT, 17:66-77, 203-215
- encephalitis, 18:34-37
- epilepsia partialis continua following pertussis infection, 16:45-47
- epilepsy, spectral analysis of interictal EEG, 16:94-97

- episodic anterior drowsy theta in adults, 17:135-138
- fever in adult patients, 18:85-88
- FIRDA, (See Frontal intermittent rhythmic delta activity)
- glioblastoma, with spindle activity awake, 17:1-5
- hepatic encephalopathy, 14:86-89
- hydrocephalus, 15:22-31
- integrated delta and spindle components in mentally retarded children, 17:24-29
- Lennox-Gastaut syndrome, 17:117-126
- lithium-specific EEG changes, 18:46-53
- lymphocytic leukemia in children, 14:214-220; 16:208-212
- neonatal meningitis, 14:67-77
- neonatal seizures, 15:125-139
- neurological disease, 14:116-129
- normal waves in children, 15:92-96
- paroxysmal hyperventilation responses in the adult EEG, 17:61-65
- periodic EEG activity, 15:32-39
- pial-dural arteriovenous malformation, FIRDA, 18:227-232
- porencephaly and subarachnoid cyst, 16:21-29
- post-traumatic syndrome, 15:214-221
- rage attacks and self-injurious behavior in brain-injured patients, 17:169-172
- rhythmic temporal theta, 15:145-150
- saline activation of pseudoseizures, 16: 171-176
- schizophrenia, treatment resistant, brain surface activity mapping, 18:124-135
- seizure control by mesothalamic reticular stimulation, 17:52-60
- seizures with pain, 16:195-201
- sleep in infants with congenital cerebral malformations, 17:92-104, 152-157
- suppression-burst pattern, (See Suppression-burst)
- tardive dyskinesia, 15:222-231; 17:30-35
- theta coma, (See Theta coma)
- topographic mapping, artifacts, 18:61-67
- tuberous sclerosis, 16:219-224
- West's syndrome with pyrethroid therapy treatment, 15:140-144
- Small sharp spikes**, and epilepsy, possible relationship, 18:114-119
  - further data on a controversial pattern, 15:208-213
- Sodium valproate**, (See Valproic acid)
- Somatic complaints** and, disturbed sleep, 18:81-84
- Somatosensory evoked potentials** and, acupuncture, 14:53-56
  - alcoholics, short latency, 14:8-16
  - alpha coma, 18:103-113
  - brainstem death, 16:63-71
  - carotid endarterectomy, assessment of

- variability of early scalp-components, 16:157-160
- dementias, cortical and subcortical, 18:26-33
- false negative intraoperative SEP with simultaneous bilateral stimulation, 17:6-9
- hepatic injury status, 18:15-19
- hypsarhythmia-infantile spasms in near-drowning, 18:180-186
- liver transplantation, monitoring cerebral functions, 16:192-194
- lumbar disc disease, dermatomally stimulated, 14:152-160
- meralgia paresthetica, stimulation of cutaneous femoris lateralis nerve, 14:161-163; 15:No.1,VII
- renal failure, chronic, short-latency changes during hemodialysis, 14:202-206
- spinal cord lesions, long term effects on configuration, 14:142-151, 173-185
- status epilepticus, non-convulsive, 16:33-38
- thoracic outlet syndrome, diagnostic importance, 17:112-116
- tibial, peroneal and sural nerve stimulation using 4 different montages, 16:149-156
- trigeminal nerve stimulation, 14:61-66
- Somatosensory spike epilepsy** and, benign infantile epilepsies, 16:39-44
- Spectral analysis**, and acute lymphocytic leukemia of children, 16:208-212
- children with end stage renal disease before and after haemodialysis, 17:195-202
- delta and spindle components during sleep in infants with congenital cerebral malformation, 17:152-157
- EEG pattern during electroconvulsive therapy, 17:66-77, 203-215
- interictal EEG in epilepsy, 16:94-97
- lorazepam and methylphenidate in normal subjects, 17:176-180
- phenytoin infusion in seizure patients, 15:173-177
- Speech disorder** and, stuttering and seizures, adult onset, 17:142-145
- Spells**, and 24 hour ambulatory EEG monitoring in pediatrics, 15:164-166
- (See also Seizures)
- Sphenoidal EEG recording**, and complex partial seizures, 17:139-141
- Spike and wave activity** and, absence status with eye-closure, 16:111-118
- benign infantile epilepsies, occipital, 16:39-44
- coma, with other patterns, 18:74-80
- complex partial status epilepticus, 14:225-232
- ECT, 17:203-215
- electrode recording systems, 15:71-82
- grand mal discharge, 15:155-158
- hypsarhythmia, natural history, 16:128-130
- Lennox-Gastaut syndrome, 15:61-68, 17:117-126; 18:187-190, 191-200
- Lennox-Gastaut syndrome with status epilepticus, 18:89-92
- mid-line activity in adult EEGs, 16:54-59
- neonatal status epilepticus, 15:197-201
- photosensitive epilepsy, 15:47-52
- post-traumatic syndrome, 15:214-221
- seizure control by mesothalamic reticular stimulation, 17:52-60
- tuberous sclerosis, 16:219-224
- unusual patterns in children, 15:92-96
- West's syndrome evolution, 15:61-68
- writing epilepsy, 17:21-23
- Spikes** and, ambulatory EEG in community hospital, analysis of patient outcome, 16:161-166
- anoxic encephalopathy, 17:187-194
- benign focal epilepsy of childhood, individual and intrafamilial multifocality of spikes, 18:169-172
- benign infantile epilepsies, evoked, 16:39-44
- burst suppression in infants, following asphyxia, 14:214-220
- Cannon and Rosenbluth's law of denervation, 17:No. 4, VII, VIII
- cerebral infarction, comparative EEG and CT scan study, 18:20-25
- clinical EEG variables suggesting extra-pyramidal side effect risk, 18:173-179
- complex partial status epilepticus, 14:225-232
- controversial patterns in children, 15:92-96
- Creutzfeldt-Jacob disease, 18:211-217
- depth studies in Lennox-Gastaut syndrome, 18:191-200
- ECT, 17:203-215
- EEG vs. CT in neurological diagnosis, 14:116-129
- electrocorticograms, 15:83-91
- evoked sharp wave excitability, 17:127-134
- evoked spike epilepsy in childhood, 14:221-224
- glioma of the sylvian region with post-ictal gustatory hallucination, 17:89-91
- grand mal discharge, 15:155-158
- herpes simplex virus encephalitis, survival in an elderly man, PLEDs, 18:2-9
- hydrocephalus, 15:22-31
- hypertensive encephalopathy, 15:53-60
- hypsarhythmia, (See Hypsarhythmia)
- interictal EEG prediction of response to drug monotherapy, 16:131-135
- Lafora disease, 16:167-170, 202-207
- Lennox-Gastaut syndrome, runs, grand mal discharge, 17:117-126
- lymphocytic leukemia, acute, 14:214-220
- mid-line, 15:232-236; 16:54-59
- multiple sclerosis, PLEDs, 18:38-40
- near-drowning, with hypsarhythmia, 18:180-186

- neonates, 15:133-139, 197-201
- Neuro-Behcet syndrome, 15:122-124
- neurophysiological basis of epilepsy, 18:163
- occipital evoked spike epilepsy in childhood, 14:221-224
- paralysis, ictal, status epilepticus, 18:12-14
- parietal epilepsy with evoked and spontaneous spikes in siblings, 17:159-161
- paroxysmal dystonic choreoathetosis, 15:151-154
- PLEDs, (See Periodic lateralized epileptiform discharges)
- post-traumatic syndrome, 15:214-221
- psychomotor type focus, 14:No.3,V
- Rett's syndrome with Lennox-Gastaut pattern, 18:187-190
- seizure control by mesothalamic reticular stimulation, 17:52-60
- small sharp spikes, controversial pattern, 15:208-213
- small sharp spikes, possible relationship to epilepsy, 18:114-119
- somatosensory epilepsy, 14: 221-224
- sphenoidal EEG recording in complex partial seizures, 17:139-141
- status epilepticus (See Status epilepticus)
- stuttering and seizures, adult onset, 17: 142-145
- subacute spongiform encephalopathy, periodic, 18:147-158
- suicidal behavior, 14:20-26
- suppression-burst pattern with alpha-theta pattern coma, 17:82-88
- tardive dyskinesia, 15:222-231
- temporal minor slow and sharp activity and cerebrovascular disorder, 18:201-210
- tuberous sclerosis, 16:219-224
- (See also Spike and wave)
- Spinal cord lesions** and, alcoholics with evoked potentials 14:8-16
- long-term effects on configuration of SEP, alterations in waveform characteristics, 14:142-151
- long-term effects on configuration of SEP, changes in components produced by afferent interaction, 14:173-185
- somatosensory evoked potentials with false negative interpretation, 17:6-9
- Spindle coma** and, benzodiazepine toxicity, case report, 16:213-218
- coexisting with alpha and theta pattern coma, 18:74-80
- (See also Spindles, sleep)
- Spindles, sleep**, and alpha coma and clinical and EP studies, 18:103-113
- benzodiazepine toxicity, case reports, 16:213-218
- coma, 14:47-52, 207-213; 16:213-218; 18: 34-37, 74-80, 103-113
- glioblastoma, case with waking spindle

- activity, 17:1-5
- infants with congenital cerebral malformation, 17:92-104, 152-157
- mentally retarded children, cyclic variation of integrated spindle components during nocturnal sleep, 15:40-46
- mentally retarded children, integrated delta and spindle components, 17:24-29
- sleep-waking disturbances in vascular brain stem lesions, 14:35-46
- viral encephalitis coma, 18:34-37
- Spongiform encephalopathy**, subacute, and, periodic paroxysmal activities, clinical evolution and serial EEG findings in 20 cases, 18:147-158
- Status epilepticus** and, absence status appearing on eye-closure, 16:111-118
- complex partial, 2 cases, 14:225-232
- control of absence status by mesothalamic reticular stimulation, 17:52-60
- epilepsia partialis continua following pertussis infection, 16:45-47
- hypertensive encephalopathy, 15:53-60
- Lennox-Gastaut syndrome, 17:117-126
- Lennox-Gastaut syndrome, clinical characteristics and management, 18:89-92
- neonatal, clinical aspects, 15:193-196
- neonatal, EEG aspects, 15:197-210
- Neuro-Behcet syndrome, 15:122-124
- non-convulsive, evoked potentials, 16:33-38
- paralysis, ictal, EEG confirmation of its epileptic nature, 18:12-14
- somato-inhibitory, EEG confirmation of a rare condition, 18:12-14
- subclinical delta status in newborn, an unfavorable prognostic sign, 15:125-131
- termination and suppression of inter-ictal bursts with phenytoin, case report, absence status, 17:173-175
- Stereo-EEG**, (See Depth recording)
- Stimulation, electrical** and, seizure control by mesothalamic reticular stimulation, 17:52-60
- Stuttering**, and seizures, adult onset, 17:142-145
- Subacute immunosuppressive measles encephalitis** (SIME), in children with acute lymphocytic leukemia, clinical EEG and CT features, 14:214-220
- Subacute sclerosing panencephalitis** (SSPE) and, EEG vs. CT in diagnosis, 14:116-129
- periodic patterns, 15:32-39
- resemblance to measles encephalitis, 14:214-220
- Subacute spongiform encephalopathy** and, periodic paroxysmal activities, clinical evolution and serial EEG findings in 20 cases, 18:147-158
- Succinylcholine chloride** (Anectine), and attenuation of EMG with electrocerebral inactivity, 16:72-76
- Suicide risk**, and patients with EEG dysrhyth-

- mias taking oral contraceptives, a speculative empirical note, 16:88-93
- relationship to EEG, qualitative considerations and report on series of completed suicides, 14:20-26

**Suppression-burst pattern** and, anoxic encephalopathy, coexistence with periodic EEG pattern, 17:187-194

- extreme stereotypy in the pattern, 17:162-168
- full term infants following asphyxia, reactive, 14:138-141
- natural history of hypsarhythmia, 16:128-130
- neonatal seizures, 15:133-139
- periodic EEG activity correlations, 15:32-39
- spikes with alpha-theta pattern coma, 17:82-88
- subacute spongiform encephalopathy, 18:147-158

**Supranuclear palsy, progressive**, and, electrophysiologic validation of cortical and subcortical dementia, 18:26-33

**Sural nerve**, and SEP stimulation, 16:149-156

**Surgery**, and electrocorticography during seizure surgery, 15:83-91

- posterior fossa, with ABR monitoring, 16:83-87
- spinal cord tumor, false negative SEP with simultaneous bilateral stimulation, 17:6-9
- (See also Depth recording)

**Syncope** and, hyperventilation responses in the adult EEG, 17:61-65

- post-traumatic syndrome, 15:214-221
- success of EEG in confirming epilepsy, 16:98-103
- (See also Pseudoseizures)

**Tardive dyskinesia** and, EEG, 15:222-225, 226-231

- 5 year follow-up of EEG and neuropsychological correlates, 14:186-201
- psychiatric inpatients, antipsychotic induced, 17:30-35

**Technique, EEG**, and, APEEG system, 15:1-21

- historical notes on bipolar and monopolar recordings, 14:No.2,V; No.3,VII,VIII
- Neo-EEG system, 15:71-82
- standards for EEG instruments, 15:No.3,VI

**Telephone polysomnogram**, in subject's home, 14:96-101

**Temporal lobe status epilepticus**, 2 case reports, 14:225-232

**Temporal localization** and, benign focal epilepsy of childhood, spikes, 18:169-172

- burst suppression pattern, slowing, 17:162-168
- complex partial seizures, sphenoidal recording, spikes, 17:139-141
- complex partial status epilepticus, 14:225-232
- ECT, analysis of spectral energy, 17:66-77
- ECT, methohexital, Innovar and ketamine

anesthesias, slowing, 17:203-215

- electrocorticograms, spikes, 15:83-91
- electrode placement systems, 15:1-21
- herpes simplex virus encephalitis, survival in an elderly man, periodic sharp complexes, 18:2-9
- hypnogenic paroxysmal dystonia seizures, 16:136-142
- hypsarhythmia, natural history, 16:128-130
- law of denervation, spikes, 17:No. 4, VII
- Lennox-Gastaut syndrome, later development, 17:117-126
- minor slow and sharp activity and cerebrovascular disorder, 18:201-210
- psychomotor type spike focus, 14:No.3,V
- rhythmic temporal theta, 15:145-150
- seizures with auditory event-related potentials, temporal lobe, 17:10-13
- small sharp spikes, controversial pattern, 15:208-213
- small sharp spikes, possible relationship to epilepsy, 18:114-119
- status epilepticus, 14:225-232
- stuttering and seizures, adult onset, spikes, 17:142-145
- very fast beta, 15:237-241

**Thalamus** and, coexistence of alpha, theta and spindle coma, 18:74-80

- depth stimulation in seizures with pain, 16:195-201
- evoked potential and clinical studies in alpha coma, 18:103-113
- evoked potentials in alcoholics, 14:8-16
- glioblastoma with spindle activity in the waking EEG, 17:1-5
- neurophysiologic mechanisms in ECT seizures, 17:203-215
- seizure control by mesothalamic reticular stimulation, 17:52-60

**Theta activity** and, Alzheimer's disease, 18:164

- brainstem death, 16:63-71
- centenarians, 17:146-151
- cerebral fat embolism, 14:27-34
- cerebellar syndrome, cerebral electrical field mapping, 18:142-146
- children with end stage renal disease before and after haemodialysis, 17:195-202
- coma, 14:47-52, 207-213; 15:32-39, 116-121; 16:No.3,VII; 17:82-88, 187-194; 18:54-60, 74-80
- ECT, analysis of spectral energy, 17:66-77
- ECT with spectral energy distributions and methohexital, Innovar and ketamine anesthesias, 17:203-215
- episodic anterior drowsy, theta in adults, 17:135-138
- fever in adult patients, 18:85-88
- post-traumatic syndrome, 15:214-221
- rhythmical temporal theta, a distinctive EEG entity, 15:145-150

- schizophrenia, treatment resistant, brain surface activity mapping, 18:124-135
- suppression-burst pattern with epileptiform discharges, 17:82-88
- tardive dyskinesia, 17:30-35
- temporal minor slow and sharp activity and cerebrovascular disorder, 18:201-210
- topographic mapping, artifacts, 18:61-67
- (See also Slow activity)
- Theta coma** and, anoxic encephalopathy, with coexistence of ectopic rhythms and periodic EEG pattern, 17:187-194
- coexisting with alpha pattern coma and spindle coma, 18:74-80
- mixed frequencies, correlations, 14:47-52, 207-213
- pattern occurring in younger adults, 18:54-60
- periodic activity, 15:32-39
- regarding initial reports, 16:No.3,VII
- suppression burst pattern, 17:82-88
- variant of alpha pattern coma, 15:116-121
- Thoracic outlet syndrome**, and diagnostic importance of SEP, 17:112-116
- Thought processes** and, FIRDA, 16:1-10
- (See also Cognitive dysfunction)
- Three per second spike and wave discharges**, and neurophysiologic mechanisms, 17:203-215
- (See also Petit mal)
- Tibial nerve** and, SEP stimulation, 16:149-156
- effect of spinal cord lesions on SEPs, 14:142-151
- Tongue movement artifacts**, in the EEG, 14:57-59
- Tonic-clonic seizures**, (See Grand mal and Seizures)
- Topographic, mapping of EEG**, See Brain electrical activity mapping
- Tracé alternant pattern** and, burst suppression in infants following asphyxia, 14:138-141
- congenital hydrocephalus, 15:22-31
- Trauma, head**, and, childhood, EEG study, 18:10-11
- EEG vs. CT in neurological diagnosis, 14:116-129
- electrophysiological tests and post-concussion syndromes, 18:165-167
- periodic EEG activity, 15:32-39
- post-traumatic syndrome, EEG features, 15:214-221
- rhythmic temporal theta, 15:145-150
- theta coma, 15:116-121
- theta pattern coma occurring in younger adults, 18:54-60
- whiplash injury, EEG reappraisal, 18:41-45
- Trigeminal nerve stimulation**, and cortical evoked potentials in humans, 14:61-66
- Triphasic delta waves** and, cerebral fat embolism, 14:27-34
- Creutzfeldt-Jacob disease, 18:211-217
- EEG vs. CT in neurological diagnosis, 14:116-129
- generalized periodic EEG activity correlations, 15:32-39
- Tuberous sclerosis** and, long-term follow-up and longitudinal EEG study, 16:219-224
- West's syndrome, 16:122-127
- Tumor, brain**, and BAER, 15:110-115
- brainstem death, 16:63-71
- EEG vs. CT in diagnosis, 14:116-129
- FIRDA, 14:135-137
- gustatory hallucination and microspikes, with sylvian region glioma, 17:89-91
- monocular VEP, crossed asymmetry, 16:77-82
- periodic EEG abnormalities, 15:32-39
- spindle activity in the waking EEG, 17:1-5
- Tumor, spinal cord**, and false negative intra-operative somatosensory evoked potentials with simultaneous bilateral stimulation, 17:6-9
- Twenty-four hour ambulatory EEG** and, monitoring in pediatrics, 15:164-166
- post-traumatic syndrome, 15:214-221
- Two per second spike and wave discharges**, (See Petit mal variant pattern)
- Typhoid and paratyphoid vaccine**, and pyretotherapy treatment in West's syndrome, 15:140-144
- Ulnar nerve**, and thoracic outlet syndrome, diagnostic importance of SEP, 17:112-116
- Universal APEEGE system**, anatomical placement of EEG electrodes, 15:1-21
- Vaccination**, and diminishing incidence of hypsarrhythmia, 16:178-182
- Valproic acid** and, effect of different doses on serum levels of total and free diphenylhydantoin, 16:104-110
- interictal EEG prediction of response to monotherapy, 16:131-135
- Lennox-Gastaut syndrome with status epilepticus, 18:89-92
- seizure management in mentally retarded epileptics, 14:164-172
- sleep and waking disturbances in epilepsy, 18:136-141
- treatment of epilepsy during the pediatric age, 17:14-20
- writing epilepsy, 17:21-23
- Vascular disease** and, alpha coma, 15:167-172
- anterior bradyrhythmia, 16:1-10
- auditory brainstem response in surgical monitoring, 16:83-87
- brainstem death, 16:63-71
- brainstem lesions and disturbance of sleep-waking cycle, 14:35-46
- carotid endarterectomy, with SER, 16:157-160
- cerebral infarction, EEG and CT scan



- comparative study, 18:20-25
- dizziness, 15:110-115
- EEG vs. CT in diagnosis, 14:116-129
- evoked potentials and dizziness, 15:110-115
- FIRDA, 14:135-137; 16:1-10; 18:227-232
- FIRDA in pial-dural arteriovenous malformation, 18:227-232
- headache with paroxysmal hyperventilation responses, 17:61-65
- hypertensive encephalopathy and anatomoclinical evidence of posterior cerebral damage, 15:53-60
- low voltage EEG, 14:102-105
- neonatal seizures, 15:125-131, 133-139, 193-196
- pediatric cases of coma, 14:207-213
- rhythmic temporal theta, 15:145-150
- temporal minor slow and sharp activity, 18:201-210
- theta pattern coma in younger adults, 18:54-60
- visual EP with crossed asymmetry, 16:77-82
- Ventricular enlargement**, and mid-line epileptiform activity in adult EEGs, 16:54-59
- Versive seizures**, and benign infantile epilepsies, 16:39-44
- Vertex localization** and, mid-line epileptiform activity in adult EEGs, 16:54-59
- mid-line spike discharges, 15:232-236
- Vertigo** and, whiplash injury, EEG reappraisal, 18:41-45
- (See also Dizziness)
- Very fast beta activity** and, clinical correlates, 15:237-241
- Vestibular dysfunction**, and BAEP, 15:110-115
- Viral infections**, and diminishing incidence of hypsarrhythmia, 16:178-182
- (See also Infectious disease)
- Visual disorder** and, benign infantile epilepsies, 16:39-44
- Cannon and Rosenblueth's law of denervation, 17:No. 4, VII
- Lafora disease, 16:167-170
- occipital evoked spike epilepsy in childhood, 14:221-224
- Visual evoked potentials (VEP)** and, alcohol abuse, pattern reversal, 15:102-109
- alcoholics, 14:17-19
- brainstem death, 16:63-71
- cerebellar syndrome, cerebral electrical field mapping, 18:142-146
- crossed asymmetry, normal and abnormal, 16:77-82
- dementias, cortical and subcortical, 18:26-33
- drowning outcome, predictors of survival in young children, 16:183-191
- gender and age changes, latency, 18:159-162
- hepatic injury status, 18:15-19
- hydrocephalus, 15:22-31
- hypsarrhythmia-infantile spasms in near-drowning, 18:180-186
- myelin disturbance in children treated for acute lymphoblastic leukemia, latency, 18:68-73, No. 3, VI-VII
- neonatal meningitis, 14:67-77
- nitrous oxide abuse and recovery, 16:143-148
- psychiatric patients, 15:178-183
- renal hemodialysis, 15:97-101
- status epilepticus, non-convulsive, 16:33-38
- Visual seizures** and, Lafora disease, 16:167-170
- (See also Occipital seizures)
- Voltage**, low, and EEG overview, 14:102-105
- coma, pediatric cases, 14:207-213
- delirium tremens, 16:30-32
- hydrocephalus, 15:22-31
- neonatal meningitis, 14:67-77
- neurological disorders, 14:116-129
- nocturnal sleep in infants with congenital malformation, 17:92-104
- (See also Electrocerebral inactivity)
- Waking EEG**, and, spindle activity with hemispheric glioblastoma, 17:1-5
- West's syndrome** and, diminishing incidence of hypsarrhythmia, 16:178-182
- evolving into Lennox-Gastaut syndrome, 15:61-68
- infection as a possible etiology of hypsarrhythmia, 17:78-81
- natural history of hypsarrhythmia, 16:128-130
- pyretotherapy as treatment, 15:140-144
- treatment, 16:122-127
- valproic acid treatment, 17:14-20
- (See also Hypsarrhythmia)
- Whiplash injuries**, and EEG features, 15:214-221
- EEG reappraisal, 18:41-45
- Wicket spikes** and, temporal minor slow and sharp activity and cerebrovascular disorder, 18:201-210
- Wilson's disease**, and SEP monitoring cerebral functions following liver transplantation, 16:192-194
- Writing epilepsy** and, two cases, seizures precipitated by writing and typing, 17:21-23



